

ISSN 0253-4924

# **FOOD TECHNOLOGY Abstracts**

**Vol. 27 No. 10 October 1992**



Central Food Technological Research Institute, Mysore  
National Information System for Science and Technology  
Department of Scientific and Industrial Research, New Delhi.



## **NATIONAL INFORMATION CENTRE FOR FOOD SCIENCE AND TECHNOLOGY**

**CFTRI, MYSORE - 570 013**

The National Information Centre for Food Science and Technology (NICFOS), is a discipline oriented information service in Food Science, Technology and Nutrition. Set up in October 1977 at the Central Food Technological Research Institute (CFTRI), Mysore, it is one of the Sectoral Information Centres under the NISSAT of the Govt. of India, Dept. of Scientific and Industrial Research.

NICFOS's services are generated by a band of professional specialists and backed up by reprographic, micrographic, printing and computer facility. Its services include publication of R & D and industry oriented current awareness services, answering technical inquiries, reprography and training.

### **NICFOS BASE**

The well-equipped Library of CFTRI with its collection going back to 1950 function as a clearing house for information on all aspects of food area. With its large collection of books, monographs, conference proceedings, bound volumes of periodicals, standards, patents, reports, theses, microforms and 600 current periodicals it can provide you with the latest information on any subject area of food. The library also serves as a training centre in different aspects of information handling and use besides guiding in setting up of new libraries in the area of food science and technology.

### **PHOTOCOPY (XEROX) OF PUBLISHED SCIENTIFIC/TECHNICAL ARTICLES**

The National Information Centre for Food Science and Technology (NICFOS) at the Central Food Technological Research Institute, Mysore, has got a good collection of scientific and other periodicals collected over the years in the area of Biological Sciences. If any article is needed for reference work, we will supply one copy from our wealth of collections. The cost of providing the copies is Rs.2/- per page (minimum charge Rs.10.00). Please take advantage of this facility to overcome your problem in getting original articles.

All correspondences regarding these services should be addressed to:

*The Area Co-ordinator*

*FOSTIS, CFTRI,*

*Mysore 570 013,*

*Karnataka, India.*



# **FOOD TECHNOLOGY ABSTRACTS**

**Vol. 27 No. 10**

**October 1992**

**National Information Centre For Food Science And Technology  
Central Food Technological Research Institute,  
Mysore - 570 013, India**

**Compiled and Edited by**

B. Vasu

C. S. Anita

Geetha Seetharam

**Abstractors to FTA**

AS	Author's Summary
BV	B. Vasu
CSA	C. S. Anita
GS	Geetha Seetharam
KAR	K. A. Ranganath
SD	S. Dhanaraj
SRA	S. R. Ananthnarayan
VKR	V. Krishnaswamy Rao

**Computerisation and Database Creation**

P. Manilal

C. S. Anita

B. Vasu

S. R. Ananthnarayan

Annual Subscription : Rs.250.00. US \$ 85.00. Single Copy : Rs.25.00



## CONTENTS

	Page No
General	... 495
Food Processing	... -
Food Packaging	... 495
Food Engineering and Equipment	... 495
Energy in Food Processing	... -
Food Chemistry and Analysis	... 496
Food Microbiology and Hygiene	... 497
Biotechnology	... -
Tissue Culture	... -
Food Additives	... 499
Cereals	... 500
Millets	... 501
Pulses	... 501
Oilseeds and Nuts	... 502
Tubers and Vegetables	... 505
Fruits	... 508
Confectionery, Starch and Sugar	... 511
Bakery products	... 514
Milk and Dairy products	... 515
Meat and Poultry	... 518
Seafoods	... 522
Protein Foods	... 525
Alcoholic and Non-alcoholic Beverages	... 525
Fats and oils	... 526
Spices and Condiments	... 529
Sensory Evaluation	... 530
Food Storage	... -
Infestation Control and Pesticides	... -
Biochemistry and Nutrition	... -
Toxicology	... 530
Food Laws and Regulations	... -
Author Index	... 531
Subject Index	... 537



## ABBREVIATIONS

A	ampere	g	gram	qt	quart
AAS	atomic absorption Spectrometry	GC	gas chromatography	R	rontgen
abstr.	abstract	gn	gravity	rad	rad or radian
ad lib.	ad libitum	gal	gallon	ref.	reference(s)
ADP	adenosine diphosphate	gf	gram-force	rev/min	revolutions per minute
Anon.	Anonymous	GLC	gas-liquid chromatography	RH	relative humidity
AOAC	Association of Official Analytical Chemists	h	hour	RNA	ribonucleic acid(s)
approx.	approximately	ha	hectare	S.	south, Southern, etc.
atm	atmosphere	HDPE	high density polyethylene	s.d.	standard deviation
ATP	adenosine triphosphate	hl	hectolitre [100 l]	SDS	sodium dedecylsulphate
$a_w$	water activity	hp	horse power	s.e.	standard error
BHA	butylated hydroxyanisole	HPLC	high performance/pressure liquid chromatography	s	second [time]
BHT	butylated hydroxytoluene	HTST	high temperature short time	SNF	solids-not-fat
BOD	biological oxygen demand	Hz	hertz [frequency cycle/s]	sp., spp.	species
b.p.	boiling point	in	inch	sp.gr.	specific gravity
Btu	British thermal unit	IR	infrared	summ.	summary
c-	centi- [as in cm, cm <sup>2</sup> , cm <sup>3</sup> ]	IU	international unit	Suppl.	Supplement
cal	calorie	J	joule	t	metric tonne
cd	candela	k-	kilo- [as in kcal, kg]	temp.	temperature
Ci	curie	K	Kelvin	TLC	thin layer chromatography
CMC	carboxymethyl cellulose	l	litre	TS	total solids
COD	chemical oxygen demand	lb	pound	UHT	ultra-high temperature
coeff.	coefficient	lb	pound-force	UV	ultraviolet
conc.	concentrated	LDPE	low density polyethylene	V	volt
concn.	concentration	m-	milli- [as in mg, ml, mm]	var.	variety
cv.	cultivar	m-equiv	milli-equivalent	vol.	volume
cwt	hundredweight	m	molar concentration	v/v	volume/volume
d-	deci-	M-	mega- [as in Mrad]	w	watt
DE	dextrose equivalent	max.	maximum	W.	West, Western, etc.
detn.	determination	min	minute [time]	WHO	World Health Organization
DFD	dark firm dry	min.	minimum	w/v	weight/volume
diam.	diameter	mol	mole	wk	week
dil.	dilute	mol.wt	molecular weight	wt.	weight
DM	dry matter, Deutsche Mark	m.p.	melting point	yd	yard
DNA	deoxyribonucleic acid(s)	MPN	most probable number	yr	year
dyn	dyne	MS	mass-spectrometry	$\mu$	micro-[as in g, m]
E.	East, Eastern, etc	n-	nano-[10 <sup>-9</sup> , as in nm]	%:	per centum
ECD.	electron capture detection	N	Newton [kg m/s <sup>2</sup> ]	>	greater than
EDTA	ethylenediaminetetra acetic acid	N.	North, Northern, normal concentration	>=	greater than or equal to;
Eh	oxidation-reduction potential	NMR	nuclear magnetic resonance	<	less than
ELISA	enzyme-linked immunosorbent assay	NPU	net protein utilization	<=	less than or equal to;
f-	femto-[10 <sup>-15</sup> , as in fCi]	oz	ounce		not greater than
°F	degree Fahrenheit	p-	pico- [10 <sup>-12</sup> , as in pCi]	Chemical symbols are used for all elements.	
FAO	Food and Agricultural Organization	P	poise	<b>ABBREVIATIONS FOR LANGUAGES</b>	
FDA	Food and Drug Administration	P	probability	Language of text	
FID	flame ionization detection	Pa	Pascal [N/m <sup>2</sup> ]	Dutch	Nl
fl oz	fluid ounce	PAGE	polyacrylamide gel electrophoresis	French	Fr
f.p.	freezing point	PER	protein efficiency ratio	German	De
ft	foot, feet	p.p.b.	parts per billion	Italian	It
		p.p.m.	parts per million	Japanese	Ja
		PSE	pale soft exudative	Norwegian	No
		PTFE	polytetrafluorethylene	spanish	Es
		PVC	polyvinyl chloride	swedish	Sv
		PVDC	polyvinylidene chloride		



## GENERAL

1956

Venkataraman (LV). **Food prospects in India by the turn of the century.** *Current Science* 63(1): 1992: 20-25

The present production status along with the per capita available and the requirement in 2000 A.D. of the following food items have been described: cereals and pulses; oilseeds; fruits and vegetables; plantation crops especially spices, and beverages; and meat and its products including poultry and fish. The food grains losses in the field and in storage are discussed and the future approach for safe and biodegradable pesticides are indicated. The future role of biotechnology in finding solution to the food problem is stressed. The manufacture of processed food in India is only 1.0% and it is strongly recommended to increase this level. KAR

1957

Gehlawat (JK). **Wealth from effluents of agro-based industries.** *Proceedings of the 53rd Annual Convention of the Sugar Technologists' Association of India* 1-15: 1991

Effluents should not be considered as waste materials but with suitable processing could be converted into useful materials. The paper covers aspects like general methods of processing effluents; recovery of constituents from effluent streams; methods of isolation, concn. and purification of components; physical, chemical and biochemical methods in conversions; effluents used for enhanced regeneration of raw materials; and the use of effluents in biotechnological process industries. GS

## FOOD PROCESSING

Nil

## FOOD PACKAGING

### Packaging materials

1958

Kumar (KR) and Balasubrahmanyam (N). **Water vapour transmission rates of multi-layer flexible packaging materials.** *Journal of Food Science and Technology (India)* 29(4): 1992: 237-238

Water vapour transmission rate values of co-extruded films having different webs and laminates at conditions of 38 plus or minus 1 C with 90 plus or minus 2% rh and 27 plus or minus 1 C and 65 plus or minus 2% rh indicated a linear relationship between rh and WVTR for LDPE while hydrophilic polyamide containing composites indicated dependence on rh. The water vapour transmission rate of multi-layer films containing polyamide as the core layer ranged between 4.1 and 5.4 g/m<sup>2</sup>.day at 38 plus or minus 1 C and 90 plus or minus 2% rh, while monofilms of polyolefins had values between 1.9 and 3.0 g/m<sup>2</sup>.day. AS

## FOOD ENGINEERING AND EQUIPMENT

1959

Azuara (E), Beristain (CI) and Garcia (HS). **Development of a mathematical model to predict kinetics of osmotic dehydration.** *Journal of Food Science and Technology (India)* 29(4): 1992: 239-242

A mathematical model based on mass balances to study the kinetics for osmotic dehydration was developed. The equation is useful for any geometrical configuration and was tested using published data on apple, beef and pineapple. In all cases of relatively short time exp., good predictions were obtained for long drying times until equilibrium was reached. AS

1960

Wang (J), Wolfe (RR) and Hayakawa (K-I). **Thermal process lethality variability in conduction-heated foods.** *Journal of Food Science* 56(5): 1991: 1424-1428

No single method is available to estimate the variability of thermal process lethality ( $W_{Fp}$ ) as a function of all thermal process parameters which significantly influence  $W_{Fp}$ . Therefore, work was conducted to develop such a method. This was accomplished through screening and regression simulations/analyses. From 19 independent parameters, 12 significant ones were selected through the analysis. The selected parameters included means of slope index of heating curve ( $f_h$ ), heating medium temp., and slope index of thermal death time curve ( $z$ ), and the coeff. variations of  $f_h$  and  $z$ . A polynomial regression equation was developed for estimating  $W_{Fp}$  as a function of all parameters found significant through the regression simulation/analysis. The equation was validated experimentally. AS



1961

Tsukada (T), Sakai (N) and Hayakawa (K-I). **Computerized model for strain-stress analysis of food undergoing simultaneous heat and mass transfer.** *Journal of Food Science* 56(5): 1991: 1438-1445

A computerized method was developed for simulating transient state heat and moisture transfer and stress distribution in axisymmetric food undergoing drying process. For this development, a simultaneous heat and moisture transfer model was coupled with the virtual work principle applicable to a body undergoing volumetric changes. The developed method was verified by comparing experimental results with those predicted. Experimental results were obtained by drying cylindrically formed samples of hydrated starch granules. They included central temp., average concn. histories and photographically observed internal crack formations. AS

1962

Lee (JE) and Singh (RK). **Scraped surface heat exchanger orientation influence on particle residence time distributions.** *Journal of Food Science* 56(5): 1991: 1446-1447

Four levels of particle concn. for an "Interaction" exp. and three levels of carboxymethyl cellulose concn., flow rate, mutator speed and particle size were used for a "Tracer Only" exp. to determine particle residence times in horizontal and vertical scraped surface heat exchangers (SSHEs). Min. normalized particle residence time (NPRT) was not significantly affected by SSHE orientation in either type in the exp. However, max. NPRT was significantly higher with vertical SSHE except in the "Tracer Only" exp. Average particles travelled faster than average bulk fluid with less deviation in horizontal SSHE whereas mean particle velocity was less than average bulk fluid velocity and had greater deviation in vertical SSHE. AS

1963

Vyas (HK), Shah (CM) and Shah (US). **Different forms of corrosion in stainless steels and preventive measures.** *Indian Dairyman* 44(4): 1992: 193-202

In stainless steel, general corrosion and localised corrosion are commonly observed. The basic requirement for corrosion to take place in stainless steel material, the characteristics of general corrosion and its prevention are described. In localised corrosion, pitting, crevice corrosion, erosion corrosion, cavitation corrosion, intergranular corrosion, stress corrosion cracking are different types, the characteristics of each one is

described along with the preventive measures. Galvanic corrosion, which happens between two dissimilar metals is also described and the preventive measures are listed. GS

## ENERGY IN FOOD PROCESSING

Nil

## FOOD CHEMISTRY AND ANALYSIS

### Chemistry

1964

Gopala Krishna (AG). **Influence of water activity on autoxidation of methyl linoleate during storage.** *Journal of Food Science and Technology (India)* 29(4): 1992: 252-253

Peroxide and total carbonyl values showed no noticeable difference at 0.02 and 0.91  $a_w$ . Anisidine value and polymer content were appreciably higher at  $a_w = 0.91$ . The results showed that the development of hydroperoxides and their subsequent secondary reactions were influenced to a considerable extent at high  $a_w$ . AS

1965

Mulvihill (DM), Rector (D) and Kinsella (JE). **Mercaptoethanol, N-ethylmaleimide, propylene glycol and urea effects on rheological properties of thermally induced  $\beta$ -lactoglobulin gels at alkaline pH.** *Journal of Food Science* 56(5): 1991: 1338-1341

1966

Graham (HD). **Isolation of gellan gum from foods by use of monovalent cations.** *Journal of Food Science* 56(5): 1991: 1342-1346

Gellan gum added to foods was recovered by precipitation with 2M NaCl. The precipitate was trapped on glass wool, washed with 1M salt sol. and eluted with boiling 40%  $H_2SO_4$  or boiling  $2 \times 10^{-4}N$  NaOH. Elution with the dilute NaOH was as effective as use of 40%  $H_2SO_4$ , except with dog food, mayonnaise and salad dressing. The hydrocolloid content of the eluate was determined by the carbazole test. Recoveries from milk, beverages, salad dressing and other foods ranged from 75 - 94% (reproducibility 2-4%). The preferred cations for precipitation were  $Na^+ > K^+ > Li^+ > NH_4^+$ . For the anions, the order was  $CO_3^- > SO_4^- > Cl^-$ . AS



1967

Hernandez (E) and Baker (RA). **Candelilla wax emulsion, preparation and stability.** *Journal of Food Science* 56(5): 1991: 1382-1383, 1387

Wax in water emulsions were prepared with candelilla wax using a modified conventional method of high pressure homogenization for foods. The resulting emulsions were compared to coconut oil-in-water emulsions. A model based on spectra absorbance analysis was used to evaluate creaming rates of wax-in-water emulsions as well as coconut oil-in-water emulsions, prepared at different homogenization pressures. Creaming rates for wax-in-water emulsion were lower than for oil-in-water emulsion. Wax-in-water emulsions prepared at 4100 kPa produced suspended globules of 1.89  $\mu\text{m}$  in size; emulsions prepared at higher homogenization pressures produced smaller average particle sizes, i.e., 0.66  $\mu\text{m}$  at 27560 kPa. AS

1968

Arora (DK), Hansen (AP) and Armagost (MS). **Sorption of flavour compounds by low density polyethylene film.** *Journal of Food Science* 56(5): 1991: 1421-1423

The interactions of aldehydes, methyl ketones, methyl esters, S compounds and alkylpyrazines with LDPE film were investigated. LDPE film, suspended on a stainless steel support, was immersed in simulated milk ultrafiltrate containing a flavour compound. After shaking the glass bottles containing the sol. and film samples for 24 h at room temp. (24 plus or minus 2 C), the flavour compound was extracted from both the LDPE samples and the  $\mu\text{F}$ , and quantitated using GC. The sorption was: aldehydes (C7 - C10) 11% to 63%, methyl ketones (C7 - C10) 1.5% to 43.0%, methyl esters (C7 - C9) 6.0% to 42.0%, and S compounds 8.5% to 21%. Insignificant levels of alkylpyrazines were sorbed. For all flavour compound classes, as carbon chain length increased, the quantity of sorbed flavour compounds increased. AS

## Chemistry (Analytical)

1969

Prabhakara Raju (RV) and Raghava Naidu (R). **A new spectrophotometric method for the determination of fenvalerate (synthetic pyrethroid).** *Journal of Food Science and Technology (India)* 29(4): 1992: 260-261

A new spectrophotometric method, developed for the detn. of fenvalerate (cyano (3-phenoxyphenyl) methyl 4-chloro- $\alpha$ -(1-methyl ethyl) benzene acetate (51630-58-1), involves the hydrolysis of fenvalerate

under alkaline conditions and the coupling of liberated aldehyde with 2,4-dinitrophenylhydrazine to yield a chromophore ( $\lambda_{\text{max}} = 465 \text{ nm}$ ). Beer's law is obeyed in the range 0.1 - 3.0 p.p.m. The method is sensitive (Sandell sensitivity = 0.0048) and it could be successfully used for the detn. of fenvalerate in water sources and formulations. AS

1970

Sapp (RE) and Davidson (SD). **Microwave digestion of multi-component foods for sodium analysis by atomic absorption spectrometry.** *Journal of Food Science* 56(5): 1991: 1412-1414

Microwave digestion of a variety of low-Na dinners was evaluated for determining Na content. Dinners containing a meat, vegetable and dessert portion were blended, digested with strong acid in the microwave oven, and read by atomic absorption. Na values were compared with those obtained by an FDA method. Also, a Na standard of known concn. was added to some samples, and the recoveries measured for two different amounts added. Reproducibility of microwave digestion was tested on replicates analyzed on the same day, on consecutive days and on consecutive wks. Of the 10 dinner types tested no matrix effects or interferences were noted, and it was concluded that the microwave digestion is a rapid means of preparing a complex matrix such as a frozen dinner for Na analysis. AS

## FOOD MICROBIOLOGY AND HYGIENE

### Microorganisms

#### Bacteria

1971

Styles (MF), Hoover (DG) and Farkas (DF). **Response of *Listeria monocytogenes* and *Vibrio parahaemolyticus* to high hydrostatic pressure.** *Journal of Food Science* 56(5): 1991: 1404-1407

*Listeria monocytogenes* Scott A and CA, were subjected at 23 C to hydrostatic pressures ranging from 2,380 to 3,400 atm and *Vibrio parahaemolyticus* T-3765-1 from 680 to 1,700 atm. For *L. monocytogenes* Scott A, pressurization in ultra-high temp.-processed (UHT) milk and raw milk appeared to provide a protective effect and lessened cell death as compared to pressurization in phosphate-buffered saline (100 mM, pH 7.0). A population of about  $10^6$  CFU/mL *L. monocytogenes* was killed by exposure to 3,400 atm within 80 min at 23 C in UHT milk. A population of about  $10^6$  CFU/mL *V. parahaemolyticus* was killed by



exposure to 1.700 atm within 10 min at 23 C in clam juice. AS

1972

Zechman (LG) and Pflug (IJ). *Bacillus stearothermophilus* spore recovery altered by media concentration and formulation. *Journal of Food Science* 56(5): 1991: 1408-1411, 1414

*Bacillus stearothermophilus* spores, subjected to a range of heating times and temp. were recovered using dextrose tryptone agar and soybean casein digest agar, with and without additives, at normal and reduced concn. Results showed that a reduction in medium concn. increased recovery of heat-stressed spores. The heated spores were inhibited by 0.5% (w/v) NaCl, based on increased recovery using 1.0 TS (1.5% tryptone and 0.5% soytone) compared to 1.0 TSN (0.15% tryptone, 0.05% soytone and 0.05% NaCl). Considering the various media tested variations of tryptic soy agar (TSA) at 0.1 of manufacturers' recommended concn. resulted in increased numbers of recovered spores. Greatest recovery was obtained using 0.1 TSA, 0.1 TSAS (TSA with starch) and 0.1 TSABS (TSA with dextrose, dipotassium phosphate and starch) with none being significantly better. SRA

#### **Lactococcus lactis**

1973

Libudzisz (Z) and Galewska (E). Citrate metabolism in *Lactococcus lactis* subsp. *lactis* var. *diacetylactis* strains. *Die Nahrung* 35(6): 1991: 611-618

The formation of diacetyl, acetoin, 2,3-butylene glycol, acetaldehyde, ethanol and lactic acid during 24 h of cultivation in milk with 0.19 and 0.5% of citrate has been studied. Depending on the strain, bacteria produced 1.5 - 1.9 mg of diacetyl, 212 - 311 mg of acetoin and 137 - 156 mg of butylene glycol in 1 l milk. An increase of the citrate concn. in milk to 0.5% resulted in an increase in the production of diacetyl from 58 to 74% and of acetoin by 2.8 - 3.7 times. The strains of distinct activity of acetoin reductase produced in these conditions 2.3 - 2.7 times as much as 2,3-butylene glycol. The recovery of citrate in the form of C<sub>4</sub>-compounds ranged from 76 to 98%, yet barely 0.18 - 0.44% in the form of diacetyl. Increased concn. of citrate in milk stimulated the production of diacetyl and acetaldehyde to the similar extent, thereby it did not result in the deterioration of organoleptic qualities of starters and milk products. Within the doses used citrate did not significantly affect growth and acidifying activity of the bacteria. AS

#### **Fungi**

#### **Mucor miehei**

1974

Bhirud (VS), Subrahmanyam (VVR) and Vaidya (SD). Influence of reaction media on esterification catalysed by *Mucor miehei* lipase. *Journal of the Oil Technologists Association of India* 23(3): 1991: 44-47

During the lipase catalysed esterification, partial removal of water from reaction media by applying vacuum, either intermittently or continuously, is essential to get high degree of conversion. High free fatty acid (FFA) containing rice bran oil can be esterified with added glycerol to produce oil with low FFA by reducing the concn. of water in the reaction media, but in presence of chlorinated solvents, product having over 55% 1-monoglycerides is obtained. SRA

1975

Bhirud (VS), Subrahmanyam (VVR) and Vaidya (SD). Alkylation of primary amines with fatty alcohols using immobilized *Mucor miehei* lipase as catalyst. *Journal of the Oil Technologists Association of India* 23(3): 1991: 47-48

Stearyl amine, which has 94.3% primary amines, after reacting with octyl-, decyl-, lauryl-, and myristyl alcohol for 96 h were lowered to 59.0, 54.4, 39.98 and 19.3% respectively and the corresponding secondary amine contents were 32.3, 38.0, 51.12 and 40.9% respectively. With myristyl alcohol, 35.0% of tertiary amine was formed while it was < 6% for the other three alcohols. Using immobilized *Mucor miehei* lipase, as catalyst, asymmetric tertiary amine could be produced as done with myristyl alcohol. Alkylation was neither affected by excess use of alcohol nor by the batch size. It is concluded that *Mucor miehei* lipase can be used for alkylation of amines, and asymmetric secondary amines of high degree purity could be produced when the products are subjected to vacuum fractional distillation. KAR

#### **BIOTECHNOLOGY**

Nil

#### **TISSUE CULTURE**

Nil



1976

Suresh (P), Ingle (VK) and Vijayalakshmi (V). **Antibacterial activity of eugenol in comparison with other antibiotics.** *Journal of Food Science and Technology (India)* 29(4); 1992: 254-256

The antibacterial activity of eugenol was tested and compared with that of various antibiotics of known concn. against several Gram positive and Gram negative microorganisms. The results showed that eugenol possessed property of inhibiting the growth of microorganisms *in vitro* at a concn. of 5 g per assay. It was found to be sensitive against organisms like *Escherichia coli*, *Enterobacter sakazaki* and *Klebsiella pneumoniae* which were resistant to antibiotics like Ampicillin, Erythromycin and Sulphamethizole. AS

## Antioxidants

1977

Economou (KD), Oreopoulou (V) and Thomopoulos (CD). **Antioxidant activity of some plant extracts of the family Labiatae.** *Journal of the American Oil Chemist's Society* 68(2); 1991: 109-113

The antioxidant activities of methanol extracts of oregano, dittany, thyme, marjoram, spearmint, lavender and basil were tested in lard stored at 75 C. The concn. of extracts in lard varied from 0.01 to 0.20%. Oregano extract was found to be the most effective in stabilizing lard, followed by thyme, dittany, marjoram and lavender extracts, in decreasing order. The induction period of lard increased with antioxidant concn. After the induction period, peroxide formation proceeded rapidly, following pseudo-zero order reaction kinetics. The rate of the reaction decreased slightly with increasing plant extract concn. Combined addition of plant extracts in lard showed a low synergistic action between thyme extract and spearmint extract. AS

1978

Kashima (M), Cha (G-S), Isoda (Y), Hirano (J) and Miyazawa (T). **The antioxidant effects of phospholipids on perilla oil.** *Journal of the American Oil Chemist's Society* 68(2); 1991: 119-122

Antioxidant effect of phospholipids on the oxidation of refined perilla oil (PO;  $\alpha$ - 18:3, 54.5%; 16:0, 7.2%; 18:0, 2.6%; 18:1, 18.6%; 18:2, 15.5%, tocopherol-free (POF) and tocopherol-enriched (POR) perilla oil were investigated by measuring wt.-gains and by the oven test at 37 C. The oxidative

stability of PO was especially increased by additions of phosphatidylethanolamine (PE) and phosphatidylserine (PS), but phosphatidylcholine (PC) scarcely showed an antioxidant effect. The oxidative stability of POF was markedly low, and none of the phospholipids (PC, PE, PS) showed an antioxidant effect on the oxidation of POF. The stability of POR was lower than that of PO regardless of its higher tocopherol contents. However, the oxidation of POR was significantly suppressed by additions of PE and PS, as was observed with PO. PC showed a small antioxidant effect on the oxidation of POR. Therefore, it seems that the antioxidant effects of phospholipids, especially of PE and PS, was due to the presence of tocopherols in the perilla oil. AS

1979

Chimi (H), Cillard (J), Cillard (P) and Rahmani (M). **Peroxyl and hydroxyl radical scavenging activity of some natural phenolic antioxidants.** *Journal of the American Oil Chemist's Society* 68(5); 1991: 307-312

The autoxidation of linoleic acid dispersed in an aqueous media and the antioxidant effect of hydroxytyrosol, oleuropein, caffeic acid and tyrosol were studied. Linoleic acid autoxidation rate was estimated by the increase of conjugated diene level and by the decrease of linoleic acid content in the samples. The phenolic compounds exhibited an antioxidant activity which increased in the order: tyrosol < caffeic acid < oleuropein < hydroxytyrosol. The analysis of the hydroperoxide isomers pointed out that hydroxytyrosol, oleuropein and caffeic acid at a concn. of  $10^{-4}$  M inhibited the formation of *trans-trans* isomers in the increasing order: caffeic acid < oleuropein < hydroxytyrosol. This inhibition could be related to the ability of phenolic compounds to scavenge peroxyl radical. Tyrosol did not inhibit the formation of *trans-trans* isomers. Phenolic compounds were degraded as a consequence of their antioxidant activity and their degradation rate was positively correlated to their antioxidant efficacy. These phenolic compounds, at a concn. of  $6 \times 10^{-3}$  M, also scavenged hydroxyl radical, with an efficiency which increased in the order: tyrosol < hydroxytyrosol < oleuropein < caffeic acid. Polar substituents at the para position, such as in caffeic acid and oleuropein, were correlated with higher hydroxyl radical quenching ability. AS

## Colourants

1980

Lauro (GJ). **A primer on natural colours.** *Cereal Foods World* 36(11); 1991: 949-953



Annatto, turmeric, beet juice, paprika, cochineal/carmine, grape skin extract and caramel are the colourants covered in this article. BV

1981

Sudhir Singh and Khanna (SK). **Toxicological evaluation of non-permitted food colours - a review.** *Indian Dairyman* 44(4); 1992: 189-192

Toxicity produced by non-permitted food colours such as Auramine (C.I. Basic yellow 2), Blue VRS (CI Feed Blue 3), Malachite green (C.I. Basic green 4), Metanil yellow (C.I. Acid yellow 36), Orange II (C.I. Acid orange 7), Rhodamine B (C.I. Food red 15) are reviewed. 25 references. GS

## CEREALS

1982

Sandberg (A-S) and Svanberg (U). **Phytate hydrolysis by phytase in cereals: Effects on in vitro estimation of iron availability.** *Journal of Food Science* 56(5); 1991: 1330-1333

Wheat bran and whole meal flours of rye and oats were soaked at optimal conditions for phytate activity (55 C, pH 5) for different time intervals. Phytate and its degradation products were determined by HPLC and related to Fe solubility under simulated physiological conditions. Small amounts of phytate ( $< 1 \mu\text{mol/g}$ ) had a strong negative effect on Fe solubility. When inositol hexa- and pentaphosphates of wheat bran and rye flour were completely hydrolysed by activating endogenous phytase, Fe solubility under simulated physiological conditions increased from 3 to 53% (wheat) and 5 to 21% (rye). Addition of wheat phytase to uncooked oatmeal increased Fe solubility from 4 to 11 and in precooked to 18%, while endogenous phytase of uncooked oatmeal had less effect on phytase digestion and Fe solubility. SRA

## Rice

1983

James (J). **Rice-husk-ash cement - A review.** *Journal of Scientific and Industrial Research* 51(5); 1992: 383-393

The use of silica from rice-husk for the production of various materials, including rice-husk ash-lime binder has gained significance. In this context, potential uses of husk, nature of silica in husk, thermal degradation of husk, crystallization of silica in husk, physicochemical properties of silica ash; reactivity of silica in ash; reaction product in ash-lime-water system; kinetics and mechanism of

ash-lime reaction; processes for producing lime-rice-husk ash cement; performance of rice-husk ash cement; hydration chemistry of rice-husk ash cement and the optimum ash/lime ratio for getting good performance of ash-lime cement have been reviewed. 107 references. KAR

1984

Semwal (AD) and Arya (SS). **Effect of spices and salt on the storage stability of precooked dehydrated rice.** *Journal of Food Science and Technology (India)* 29(4); 1992: 210-213

Effect of 19 spices (1%), NaCl (0.5 - 2%) and transition metal ions ( $\text{Fe}^{++}$ ,  $\text{Cu}^{++}$ ,  $\text{Co}^{++}$ ,  $\text{Ni}^{++}$  each at 5, 50 and 500 p.p.m.) on the storage stability of pre-cooked, dehydrated rice processed using refined sunflower oil and vanaspati was investigated. All the spices except *Tejpat* (*Cinnamomum tamala*) exhibited antioxygenic activity in dehydrated rice as well as pure sunflower oil and vanaspati. Red chilli, clove mace and nutmeg exhibited max. antioxygenic activity. *Tejpat*, on the other hand, exhibited pro-oxygenic activity in all the systems. In dehydrated rice, NaCl exhibited slight pro-oxygenic activity but  $\text{Cu}^{++}$ ,  $\text{Fe}^{++}$  and  $\text{Co}^{++}$  exhibited very strong pro-oxygenic activity. Only at 500 p.p.m. level  $\text{Ni}^{++}$  exhibited marginal antioxygenic activity. Nature of the oil used during processing also significantly influenced the stability of dehydrated rice and antioxygenic or pro-oxygenic activity of the spices and metal ions. AS

1985

Hamaker (BR), Griffin (VK) and Moldenhauer (KAK). **Potential influence of a starch granule-associated protein on cooked rice stickiness.** *Journal of Food Science* 56(5); 1991: 1327-1329, 1346

The amount of a 60 Kdal starch granule-associated protein, or waxy gene product, found in milled white rice was examined in relation to cooked rice stickiness. For 32 rice selections the correlation coeff. was similar though slightly lower for the 60 Kdal protein and stickiness ( $r = -0.85$ ,  $P < 0.01$ ) compared to amylose and stickiness ( $r = -0.87$ ,  $P < 0.01$ ). The 60 Kdal protein content correlated highly with amylose content ( $r = 0.96$ ,  $P < 0.01$ ), suggesting a more clear association between amylose and texture. The gelatinization properties of the starch granule may be influenced by the relative amount of granule-associated protein. SRA

## Wheat

1986

Syed (HM), Indoni (SB), Annapure (SG) and Bodhankar (SS). **Physico-chemical**



**characteristics of Indian wheats - a review.** *Indian Miller* 22(5): 1992: 27-31

A brief review covers physical parameters, protein content, gluten content and mineral content of the Indian wheat. 36 references. SRA

## **Wheat flour**

1987

Indrani (D) and Venkateswara Rao (G). **Effect of milling methods on the chemical, rheological and breadmaking characteristics of whole wheat flour.** *Journal of Food Science and Technology (India)* 29(4): 1992: 218-220

Whole wheat flour obtained by milling wheat in hammer, disc, stone and roller mills showed considerable variation in particle size distribution. The damaged starch content and diastatic activity ranged from 10.7 to 21.2% and from 187 to 380 mg of maltose/10g of flour respectively thereby reflecting variation in degree of severity of grinding in different mills. The dough raising capacity was also affected with the values ranging between 53.4 and 79.6%. The dough properties of flours showed considerable variation and the water absorption varied from 64.9 to 72.6%. Bread samples prepared from hammer and roller milled flours were better in quality than those from disc and stone milled flours. AS

## **MILLETS**

### **Corn**

1988

Sinha (R) and Sharada (D). **Chemical characteristics of maize grains and their relationship to roti quality.** *Journal of Food Science and Technology (India)* 29(4): 1992: 243-245

Protein, starch and sugar contents were found to decrease upon alkali treatment of maize grains in contrast to increase in amylose and ash contents. Untreated *rotis* of all var. were found significantly more acceptable than treated *rotis*. Significant correlation of protein, starch and total sugars with *roti* qualities was identified. AS

### **Corn flour**

1989

Huang (CJ) and Zayas (JF). **Phenolic acid contributions to taste characteristics of corn**

**germ protein flour products.** *Journal of Food Science* 56(5): 1991: 1308-1310, 1315

Corn germ protein flour (CGPF) was subjected to three treatments (ethanol-washing, heat treatment and acid-washing) to remove phenolic compounds. Eight phenolic acids in the extracts of CGPF samples were identified and quantified by GC. o-Coumaric, p-coumaric, and ferulic acids were the principal phenolic acids in the free and soluble ester fractions of CGPF products. Taste thresholds for the acids were 20, 48 and 90 p.p.m., respectively. Treatments generally improved sensory properties, but acid washing increased sour scores. SRA

## **Sorghum**

1990

Subramanian (V), Seetharama (N), Jabunathan (R) and Venkateswara Rao (P). **Evaluation of protein quality of sorghum (*Sorghum bicolor* (L.) Moench).** *Journal of Agricultural and Food Chemistry* 38(6): 1990: 1344-1347

## **PULSES**

1991

Idouraine (A), Yensen (SB) and Weber (CW). **Tepary bean flour, albumin and globulin fractions functional properties compared with soy protein isolate.** *Journal of Food Science* 56(5): 1991: 1316-1318, 1326

The tepary bean flour (TF) and tepary proteins had good functional properties. Teparies albumin (TA) and teparies globulin (TG) showed high protein solubility at acid and alkaline pH ranges and a low solubility at pH 4 and pH 5-6, respectively compared to soy protein isolate (SPI). Viscosity of both fractions was similar and markedly lower than that of SPI. Heating to 60 C increased viscosity in all, except in TG where it decreased. Heat coagulability and foaming properties was high in TA and TF. TG did not coagulate upon heating to 100 C. TG also formed a weaker and less stable foam and showed lower emulsion capacity than TF, TA and SPI. Oil absorption capacity was excellent to good in TA and TG fractions. SRA

1992

Vidal-Valverde (C) and Frias (J). **Legume processing effects on dietary fiber components.** *Journal of Food Science* 56(5): 1991: 1350-1352

Effects of different processes were studied on neutral detergent fiber (NDF), acid detergent fiber (ADF), hemicellulose (HMC), cellulose (CL) and lignin (LN)



content of legumes. Chickpeas, kidney beans and lentils were soaked and cooked, simulating home processing. Values were recorded on a wet basis as a guide to calculating their contribution to the diet. Normal cooking of chickpeas and kidney beans, led to higher NDF and HMC. LN increased with pressure cooking. The fiber components in lentils did not undergo any significant variation with type of cooking. SRA

### Cowpeas

1993

Okechukwu (PE), Rao (MA), Ngoddy (PO) and McWatters (KH). **Thermal processing of cowpea slurries.** *Journal of Food Science* 56(5): 1991: 1302-1307

Simple heating curves were obtained during the heating of slurries containing 15 - 30% cowpea flour. A 10% cowpea slurry at 121 C showed broken heating with two break points, one break due to gelation of protein and starch at 66 C and another due to loss of rigidity at 87 C. The  $f_h$  and  $j_h$  parameters were independent of particle size of cowpea flour. The difference in  $F_0$  at the can center and the slowest heating point were as much as 29%. SRA

### Cowpea flour

1994

Okechukwu (PE), Rao (MA), Ngoddy (PO) and McWatters (KH). **Flow behaviour and gelatinization of cowpea flour and starch dispersions.** *Journal of Food Science* 56(5): 1991: 1311-1315

The flow behaviour of a 25% cowpea slurry with 8% oil held at 70 C showed shear-thinning behaviour and an Arrhenius temp. relationship. Cowpea flour (8%) and starch (2.5%) slurries heated for < 1 min at 70 - 87 C exhibited shear-thickening while those heated longer times exhibited shear-thinning behaviour. Max. viscosities attained due to heat-induced gelatinization showed a power relationship with temp. of heating. Starch gelatinization kinetics followed a first-order equation and the temp. dependence of the rate constant followed the Arrhenius relationship with an activation energy of about 47.5 kcal/mol. Heating the slurries for > 200 min above 80 C resulted in loss of viscosity. AS

### Faba beans

1995

Zlena (HM), Youssef (MM) and El-Mahdy (AR). **Amino acid composition and some antinutritional factors of cooked faba beans (Medamnis): Effects of cooking temperature and time.** *Journal of Food Science* 56(5): 1991: 1347-1349, 1352

Easy-and hard-to-cook bean seeds were cooked by different heat treatments (100 - 125 C for 1 - 12 h). Amino acid composition, tannins, phytic acid and trypsin inhibitor activity (TIA) were determined. Almost all essential amino acids declined after cooking. Less than 10% of total tannins were decomposed during cooking, while up to 50% were leached to the cooking liquor. Retention of phytic acid in cooked beans was significantly lower than in cooked bean-liquor mixtures. Loss of phytic acid due to leaching was much higher for easy-to-cook beans than for hard-to-cook ones. Apparent retention of TIA amounted to about 50%. Optimum heat treatments were 125 C at 1 h for easy and 120 C at 2 h for hard-to-cook beans. AS

## OILSEEDS AND NUTS

### Canola

1996

Owusu-Ansah (YJ) and Marianchuk (M). **Microwave inactivation of myrosinase in canola seeds: A pilot plant study.** *Journal of Food Science* 56(5): 1991: 1372-1374, 1406

The study indicates that microwave heating could be used to completely inactivate myrosinase in whole or flaked canola seeds. The parameters for inactivation were exposure time and power. The exposure time needed for enzyme inactivation at a specific power was moisture-dependent. For constant, single processing conditions, moisture content of 10 - 16% and medium to high microwave power appeared ideal. Microwave treatment caused significant increase in the yellow colour and sulphur content of the seed oils. SRA

### Groundnuts

1997

Chlou (RY-Y), Shyu (S-L) and Tsai (C-L). **Characterization of gamma-irradiated peanut kernels stored one year under ambient and frozen conditions.** *Journal of Food Science* 56(5): 1991: 1375-1377

$\gamma$ -Irradiated (at 0, 2.5, 5.0, 10 and 20 kGy) peanuts lost germination capabilities after 1 yr storage. Peanuts irradiated with 2.5 kGy and stored at



ambient temp. contained viable molds. Peanuts oil in kernel stored at -14 C was comparatively more stable than in peanuts stored at ambient temp. Irradiation did not show any effect on oxidation of oil. Fatty acid content varied slightly with exception of linoleic and lenolenic acids which decreased with increased radiation depending on storage temp. Peanut proteins analysed by SDS-PAGE showed no obvious qualitative variation among the samples subjected to various doses of  $\gamma$ -irradiation. SRA

1998

Kim (N-K) and Hung (Y-C). **Mechanical properties and chemical composition of peanuts as affected by harvest date and maturity.** *Journal of Food Science* 56(5): 1991; 1378-1381, 1392

The effects of harvest date and maturity (by the Hull Scrape method) on the mechanical properties and the chemical composition of peanuts and the test hypothesis that peanut mesocarp colour could be used as an index to group peanuts with similar qualities were evaluated. In general, peanut kernels became harder, crisper, crunchier and more brittle at higher DAP's (days after planting) and higher maturity. Moisture content at harvest decreased, protein content remained unchanged and carbohydrates decreased with increasing DAP, harvest dates did not affect oil and ash. Total unsaturated fatty acids content increased with maturity. Qualities of peanuts in orange, brown and black maturity groups were very similar. The larger the portion of peanuts with orange, brown or black mesocarp colour, the higher would be the quality of the particular batch of raw peanuts. SRA

1999

Bhagya (S), Prakash (V) and Srinivasan (KS). **Effect of different proteolytic enzymes on the nature of subunit composition of arachins from groundnut (*Arachis hypogaea* L.).** *Indian Journal of Biochemistry and Biophysics* 29(2): 1992; 154-159

Results of this study indicated that even though the gross conformation and other physico-chemical properties of the arachins from 3 different (Spanish improved, TMV-2, and DH-3-30) var. were similar, the hydrolysis of proteins by the 3 proteolytic enzymes differed in their rate as well as extent. This can only be attributed to the nature of acidic and basic subunits of arachins in the 3 var. This is also confirmed by difference in their amino acid composition, especially in tyrosine, glutamic acid, alanine and cysteine, with possible role in the stereo-specificity of the enzymes. SRA

## Rapeseeds

## Rapeseed proteins

2000

Kroll (J). **Selected functional properties of detoxified rapeseed protein preparations effected by phytic acid.** *Die Nahrung* 35(6): 1991; 619-624

Detoxified rapeseed protein preparations with high protein content ( > 90%) produced by extraction, ultra- and diafiltration, have a very good solubility and excellent foaming properties (better than egg white). The emulsifying properties (emulsifying activity and emulsion stability) are moderate. The N-solubility profile of these preparations (albumin and globulin mixing) has been influenced by phytic acid (PA). It is possible to observe a shift of the isoelectric range to lower pH-values and parallel a decrease of solubility if the amount of PA (3, 5, 10%) is increased. The foaming properties have also been dependent on the PA content. A higher concn. of PA (3, 5, 10%) causes a decrease in the foaming activity. The results have been discussed concerning the interactions between PA and proteins. AS

## Safflowers

2001

Binder (RG), Benson (ME) and Flath (RA). **Volatile components of safflower.** *Journal of Agricultural and Food Chemistry* 38(5): 1990; 1245-1248

2002

Gupta (RK), Khanna (KR) and Banerji (R). **Variation in fatty acid composition of Indian cultivars of safflower (*Carthamus tinctorius* L.).** *Journal of the Oil Technologists Association of India* 23(3): 1991; 56-57

Fatty acid composition of 37 lines of Indian safflower is reported. The oil content of 15 samples ranged from 24 - 29%; others had > 30% oil content. All the oil samples had palmitic (16:0), oleic (18:1) and linoleic (18:2) acids, as the common fatty acids, while stearic (18:0) was found in only 15 samples. The sum of concn. of oleic and linoleic ranged from 85.42% in S-83 to 94.54% in S-99; the exception being SJ-22 (39.98%). Based on the fatty acid composition, the genotype S-48 was selected as the most promising for developing high oleic acid lines suitable for Indian conditions. SRA

## Sesame

2003

Razia Tasneem and Prakash (V). **The nature of the unhydrolysed fraction of Alpha-globulin, the major protein components of *Sesamum indicum* L. hydrolysed by Alpha-chymotrypsin.** *Indian*



*Alpha*-globulin, the high mol. wt. protein fraction from sesame (*Sesamum indicum* L.) seed, was hydrolysed by *alpha*-chymotrypsin. The hydrolysate was resolved into two fractions, the hydrolysed part and the unhydrolysed part of *alpha*-globulin using gel filtration on Sepharose 6B-100. The unhydrolysed *alpha*-globulin residue was characterized for its sedimentation coeff., subunit composition, fluorescence omission spectrum, secondary structure, and other biophysical properties. The results indicated a decrease in the size of the protein molecule upon hydrolysis to a very small extent. The effect of hydrolysis products on hydrolysis of native *alpha*-globulin as well as on a standard substrate, casein, was also investigated. The results indicated that the hydrolysis products contribute to the resistance of *alpha*-globulin to proteolysis by *alpha*-chymotrypsin to the extent of nearly 40%. AS

#### Soybeans

2004

Charjan (SKU) and Tarar (JL). **Effect of storage on germination and microflora of soybean (*Glycine max*) seed.** *Indian Journal of Agricultural Sciences* 62(7): 1992: 500-502

Soybean with 9% moisture content was stored in jute, cloth and polyethylene (700 gauge) bags (20 x 30 cm size) for 18 months at 10.6 - 43.4 C and 35.8 - 87.3% rh and observations were recorded for germination and incidence of mycoflora at 6-month intervals. Fungi incidence was max. in soybean stored in cloth and jute bags. Polyethylene bag provided good protection to soybean seeds and prevented the development of fungal colonies both quantitative and sp. wise. With the advancement of storage the incidence of various fungi increased. KAR

2005

Clark (PK) and Snyder (HE). **Hydroperoxide formation in soybean seeds during storage.** *Journal of the American Oil Chemist's Society* 68(5): 1991: 346-347

'Forrest' soybeans were stored for 2 yrs, and the extracted lipids were assayed for hydroperoxide content. The crude lipid was separated by HPLC, and 3 hydroperoxide peaks plus the triglyceride peak were measured every 2 months. There was a lag in hydroperoxide production for the first 9 months followed by a steady increase for the remaining 22 months. This method of measuring changes in lipid oxidation should be useful for

monitoring seed changes in germination and vigor during storage. AS

#### Soy products

##### Soy flour

2006

Kulkarni (SD), Wijeratne (WB) and Wei (TM). **Production of medium fat soy flour by dry-extrusion-expelling of raw soybean and its use in bread fortification.** *Journal of Food Science and Technology (India)* 29(4): 1992: 220-223

Dry heat treatment of soybean through dry-extrusion-expelling was used to produce medium fat soy flour. The flour of the cake obtained after expelling the extrudate of coarsely ground soybean contained 3.3 - 5.3% moisture and 7 - 9% oil, while trypsin inhibitor was inactivated to an extent of 35.5 - 46.8%. Soy flour thus obtained was used at 12% flour substitution level for making bread, which was acceptable. AS

2007

Mishra (HN) and Mukherjee (RK). **Storage stability of full-fat soy flour and soy-wheat flour blend.** *Journal of Food Science and Technology (India)* 29(4): 1992: 224-227

Raw full-fat soy flour (RSF), processed full-fat soy flour (FFSF) with and without antioxidant and soy-wheat flour (SWF) blend, stored at different temp. and rh, showed that the temp. and rh had pronounced effect on storage stability of the flours which were found to be more stable at lower temp. and rh ranges than at higher ones. Thermally processed and prepared FFSF had good keeping quality and could be stored safely for more than 4 months either alone or in combination with wheat flour. Addition of antioxidant had little effect in checking peroxide formation of the flour during the later stages of the storage. AS

##### Soy proteins

2008

Bernardi Don (LS), Pilosof (AMR) and Bartholomai (GB). **Enzymatic modification of soy protein concentrates by fungal and bacterial proteases.** *Journal of the American Oil Chemist's Society* 68(2): 1991: 102-105

Solubility, foaming capacity and foam stability of denatured soy protein concentrate obtained from toasted flour were improved by proteolysis with fungal or bacterial proteases. Emulsifying capacity was unchanged, but emulsion stability decreased;



bacterial protease highly improved oil absorption. Also, the bacterial protease was able to solubilize more protein and gave products which foamed more than those obtained with the fungal enzyme. However, the stabilizing properties of the bacterial modified soy protein concentrate at the air/water or oil/water interface were inferior. By limited hydrolysis up to degree of hydrolysis 10% most functional properties were improved without greatly reducing emulsion stability and water absorption. AS

2009

Kang (JII), Matsumura (Y) and Mori (T). **Characterization of texture and mechanical properties of heat-induced soy protein gels.** *Journal of the American Oil Chemist's Society* 68(5); 1991: 339-345

Heat-set gels were prepared from acid-precipitated soybean proteins at various heating temp. (80 - 100 C), protein concn. (18 - 20%), and proportions of glycinin. The gels were evaluated for mechanical parameters by means of a compression-decompression test. Gels formed at higher heating temp. and protein concn. were firm, tough and unfracturable. The elasticities of gels were similar at all protein concn. and were lower when heated at higher temp. Heating above 93 C was necessary for formation of rigid gels. The glycinin/ $\beta$ -conglycinin ratio affected the texture of the gels. Three-dimensional representation of the gels through factor analysis of instrumental data and calculation of factor scores was useful to evaluate the texture of the gels. AS

## TUBERS AND VEGETABLES

### Beet

2010

El-Makhzangy (A), Ayyad (K) and Abo-ElNile (E). **Factors affecting the production of *Saccharomyces cerevisiae* from sugar beet pulp.** *Die Nahrung* 35(6); 1991: 641-654

Factors affecting the production of *S. cerevisiae* from sugar beet pulp after acid hydrolysis were investigated. Max. yield and economic coeff. were obtained at sugar concn. of 7.15% after an incubation period of 72 h at 30 C and pH 6.0 using a mixture (1:1) of ammonium sulphate and ammonium nitrate (0.2 g/g sugar) as nitrogen source. HPLC analysis of beet pulp hydrolyzate showed that hexoses especially glucose serve as the most suitable sugars with regard to *S. cerevisiae* activity. AS

### Brassica

2011

Poonam Aggarwal and Saini (SPS). **Chlorophyll losses during preparation, canning and storage of Brassica greens.** *Journal of Food Science and Technology (India)* 29(4); 1992: 258-259

Total chlorophyll content decreased to 80% of its initial value after cooking and canning and to 40 - 50%, during 6 months storage in Brassica greens. 'RLM-240' and 'Brown sarson' retained max. chlorophyll and Chinese cabbage contained min. chlorophyll in fresh, processed as well as stored form among the 5 var. studied. Higher chlorophyll was retained under refrigerated storage. AS

### Carrots

2012

Tatsumi (Y), Watada (AE) and Wergin (WP). **Scanning electron microscopy of carrot stick surface to determine cause of white translucent appearance.** *Journal of Food Science* 56(5); 1991: 1357-1359

Carrot sticks that were prepared with a sharp culinary knife exhibited a whitish translucent appearance on the surface. This condition was not readily apparent with carrot sticks sliced with a razor sharp blade. Scanning electron microscopic examination of the translucent tissue revealed that the knife tended to shear, separate and compress the cells and tissues of the root. Dehydration of the large mass of exposed cells probably was responsible for the appearance of the whitish translucent tissue. Development of this condition is undesirable because consumers associate this with aged or nonfresh carrot sticks. AS

### Cassava

2013

Badrie (N) and Mellowes (WA). **Effect of extrusion variables on cassava extrudates.** *Journal of Food Science* 56(5); 1991: 1334-1337

Cassava flour was successfully used as feed ingredient in production of a puffed extrudate. The physicochemical characteristics of the extrudate were significantly influenced by feed moisture (FM), temp., screw speed and feed rate (FR). Optimum temp., screw speed and feed rate (FR) for extrudate expansion (2.82) was obtained at 11% FM; temp., 120 - 125 C; screw speed of 520 r.p.m. with FR 250 g/min. SRA



## Cassava flour

2014

Badrie (N) and Mellowes (WA). **Texture and microstructure of cassava (*Manihot esculenta* Crantz) flour extrudate.** *Journal of Food Science* 56(5): 1991: 1319-1322. 1364

Cassava flour (*Manihot esculenta* Crantz) was texturized by single-screw extrusion processing. Relationships between texture and microstructure as a function of extrusion variables were examined. The effect of feed moisture or screw speed was significant ( $P < 0.01$ ) on all textural parameters except springiness and energy first bite. Hardness, gumminess ( $P < 0.05$ ) fracturability and cohesiveness ( $P < 0.01$ ) decreased with increasing temp. Hardness, fracturability and firmness were lowest at screw speed 520 r.p.m. with feed rate 300g/min, 120-125 C and 11% feed moisture. Scanning electron micrographs showed structural change from a coarse cell mass at high feed moisture to a porous, thin-walled structure with decreased feed moisture. AS

## Turnips

2015

Shattuck (VI), Kakuda (Y), Shelp (BJ) and Kakuda (N). **Chemical composition of turnip roots stored or intermittently grown at low temperature.** *Journal of the American Society for Horticultural Science* 116(5): 1991: 818-822

Field and greenhouse studies were conducted to investigate the effects of low temp. on the starch, sugar, ascorbic acid, and glucosinolate (GS) concn. in turnip [*Brassica rapa* ssp. *rapifera* (Metzg.) Sinsk] roots. Field-harvested roots were stored at 0 C for 2 and 4 wks. In the greenhouse, plants were grown at 0 to 12 C for parts of 11 days before harvest. Cold-stored roots decreased in both starch and total sugar concn. (sucrose, fructose and glucose) when compared to freshly harvested roots. Greenhouse-grown plants subjected to low temp. had roots with a similar starch content but with a higher concn. of total sugars than the control. In both exp., the cold treatments induced a slight but significant increase in root sucrose concn. The ascorbic acid concn. of roots was not affected by low temp. In both the field and greenhouse studies, low temp. did not change the total concn. of the 8 major GSs identified in peeled root and peel tissues, but did alter the concn. of specific GSs. AS

## Asparagus

2010

Baxter (L) and Waters (LJr). **Quality changes in asparagus spears stored in a flow-through CA system or in consumer packages.** *Hortscience* 26(4): 1991: 399-402

Asparagus spears (*Asparagus officinalis* L.) stored 28 days at 2 C in air, a flow-through controlled-atm. (CA) system, or 14 days in polymeric film consumer packages were evaluated in respect to compositional and quality changes. CA-stored spears retained more sugars, organic acids, and soluble proteins than spears stored in air. Spears stored in vented consumer packages had a useful life of 14 days, whereas those in nonvented packages started to break down after 8 days. Spears from vented packages lost more wt. but retained more sugars and organic acids than those from nonvented packages. AS

## Potatoes

2017

Saikia (L) and Ranganna (S). **Determination of thermal process schedules for canned drumstick, okra, elephant yam and potato.** *Journal of Food Science and Technology (India)* 29(4): 1992: 203-209

Thermal process schedules have been evolved for canned drumstick, okra, elephant yam and potato using *Clostridium sporogenes* as the test organism. The values of thermal resistance (D) found using approx.  $10^3$  spores (B) were 1.75 plus or minus 0.21 times the values found using approx.  $10^4$  spores (A). The sterilisation values ( $F_0$ ) corresponding to 3.0 min (Botulinum cook), 5D of A, or a value found by integrating D values of A and B concn. of spores did not prevent spoilage of inoculated packs.  $F_0$  of 5.2 and 5.4 min (equal to 5D of B and experimental z) are considered adequate for commercial processing of drumstick and elephant yam respectively. Okra is best canned in acidified form. Potato requires an  $F_0$  of 9.7 min to overcome facultative thermophilic spoilage. A procedure for interconversion of F value corresponding to experimental z value in terms of  $F_0$  value is given. AS

## Vegetables

### Cucumbers

2018

McDonald (LC), Fleming (HP) and Daeschel (MA). **Acidification effects on microbial populations during initiation of cucumber fermentation.** *Journal of Food Science* 56(5): 1991: 1353-1356. 1359



Addition of acetic acid (0.067M) or calcium acetate (0.133 M) to the cover brine (1.94M NaCl) of cucumbers reduced naturally occurring *Enterobacteriaceae*, but not lactic acid bacteria (LAB), during the first 5 days after brining. Naturally occurring LAB were predominantly heterofermentative (greater than or equal to 80%) 1 day after brining and homofermentative (greater than or equal to 90%) on the 5th day. *Enterobacteriaceae* survived longer within cucumbers than in brine, particularly in nonacidified cucumbers. Starter cultures of *Lactobacillus plantarum* or *Pediococcus pentosaceus* decreased 90 - 99.9% during the first 10 h after addition and did not increase until about 30 h after brining. Obtaining predominant fermentation of brined cucumbers by added cultures was difficult due to presence of natural microflora on/in the cucumbers and the harsh environment of the brine (high salt, low nutrients). AS

## Leafy vegetables

### Amaranthus

2019

Bressani (R) and Garcia-Vela (LA). **Protein fractions in amaranth grain and their chemical characterization.** *Journal of Agricultural and Food Chemistry* 38(5); 1990; 1205-1209

### Pumpkins

2020

Teotia (MS). **Advance in chemistry and technology of pumpkins.** *Indian Food Packer* 46(1); 1992; 9-31

The review covers major aspects like storage, chemistry and technology of pumpkins (*Cucurbita* sp.). Under storage the effect of containers and storage conditions of fertilizers, pretreatments, surface wax and keeping quality and changes taking place during storage of pumpkins are covered. The chemistry of pumpkins and its seed covered are: varietal variation in composition, and nutritional value. Preparation of raw pumpkin for processing, blanching, dehydration, canning, pumpkin beverage, jam preserve, pickle and sauce and such other products, and the utilization of seeds are covered under the technology. 97 references. GS

### Tomatoes

2021

Larrigaudiere (C), Latche (A), Pech (JC) and Triantaphylides (C). **Relationship between stress ethylene production induced by gamma-irradiation and ripening of cherry**

**tomatoes.** *Journal of the American Society for Horticultural Science* 116(6); 1991; 1000-1003

Changes in 1-amino-cyclopropane-1-carboxylic acid (ACC) metabolism induced by  $\gamma$ -irradiation have been studied during ripening of cherry tomatoes (*Lycopersicon esculentum* L. cv. Sweet 100) treated at the mature-green stage. Irradiation caused a sharp and transient dose-dependent increase in ethylene production during the first 24 h that was associated with an increase in ACC synthase activity. The activity of ethylene-forming enzyme (EFE) was also stimulated but was never limiting. Nine days following irradiation, ACC metabolism was more active in irradiated fruits, with ACC being mainly directed to ethylene [80% at 3 kilogray (kGy; 1 Gray = 100 Rad = 1 J.kg<sup>-1</sup>)] rather than to malonyl ACC (MACC). As a consequence, fruit ripening was accelerated. For doses < 1 kGy, the time required for 50% of the fruits to reach breaker stage (the onset of climacteric ethylene production) was inversely correlated with radiation dose and the amount of stress ethylene produced during the first 24 h. At doses > 1 kGy, in spite of a continuous stimulation of stress ethylene production, no additional acceleration of ripening occurred. At 3- to 5-kGy doses, fruit ripening was impaired transiently with a fast subsequent recovery. As a result, a significant synchronization of fruit ripening (presumed to be caused by enhanced ethylene production) was observed. AS

2022

Dodds (GT), Brown (WJ) and Ludford (PM). **Surface colour changes of tomato and other solanaceous fruit during shilling.** *Journal of the American Society for Horticultural Science* 116(3); 1991; 482-490

Chilling of mature-green (MG) tomato fruit (*Lycopersicon esculentum* Mill. and related sp.) was investigated to determine the effect of chilling stress on surface colour during low-temp. storage. Colour measurements were made with a tristimulus colorimeter (L, a, b values), and data were analyzed by multivariate analysis of variance and canonical variates analysis. Changes in surface colour of MG fruit during chilling were not correlated overall with relative chilling sensitivity of cvs/lines; however, within standard and cherry types, chilling-tolerant fruit changed surface colour more during chilling than chilling-sensitive fruit when fruit were picked early in the season. Early harvests were less chilling-sensitive than late harvests. The number of hrs below 15.6 C in the 200 h before harvest was positively correlated with postharvest chilling sensitivity. A high vs. ambient rh during storage did not affect chilling-induced % change in colour. Tobacco mosaic virus resistance led to less and *Verticillium albo-atrum* Reinke & Berthier resistance



led to more chilling-induced colour change. There was no effect from resistances to *Fusarium oxysporum* Schlechtend f. sp. *lycopersici* (Sacc.) W.C. Snyder and H. N. Hans, alternaria stem canker (*Alternaria solani* Sorauer), anthracnose (*Colletotrichum coccodes* (Wallr.) S. J. Hughes), root-knot nematode (*Meloidogyne hapla* Chitwood), *Phytophthora infestans* (Mont.) deBary, or *Stemphyllium botryosum* f. sp. *lycopersici* Rotem, Cohen, and Wahl. The results show that the harvest date had an effect on chilling-induced changes in surface colour in MG fruit. AS

## FRUITS

2023

Carbonell (E), Costell (E) and Duran (L). **Fruit content influence on gel strength of strawberry and peach jams.** *Journal of Food Science* 56(5): 1991; 1384-1387

Influence of fruit content, interaction with final soluble solids and added pectin on gel strength of strawberry and peach jams was studied. Composition ranged 25 - 55%, 60 - 70 °Brix, 0.3 - 0.7% added pectin in strawberry jams and 0.1 - 0.5% in peach jams. Gel strength was measured in a Stevens LFRA texture analyzer. Regression equations showed that in strawberry jams effect of fruit content was quadratic and interaction with soluble solids was weak and negative while in peach jams the same effect was linear and the interaction was strong and positive. No interaction of fruit content with added pectin was observed, showing those effects were independent and additive. AS

### Apples

2024

Sud (G), Parmar (C) and Nayital (RK). **Effect of calcium chloride, bavistin and diphenylamine on the shelf-life of apple cv. Royal Delicious.** *Indian Food Packer* 46(1): 1992; 33-38

Apple fruits (var. Royal Delicious) after harvest were dipped in CaCl<sub>2</sub> (2 and 4%) and Bavistin (500 p.p.m.) and in diphenylamine (1000 p.p.m.) sol. for 2 min. CaCl<sub>2</sub> particularly 4% was effective in reducing the loss of fruit firmness and physiological wt., preventing the rot and maintaining the quality during 5 wk of storage at room temp. The shelf-life, which was only 2 wk under control was further extended by another 2 wk and CaCl<sub>2</sub> (4%) proved to be the best treatment. GS

2025

Lau (OL) and Yastremski (R). **Retention of quality of 'Golden Delicious' apples by controlled- and modified-atmosphere storage.** *Hortscience* 26(5): 1991; 564-566

'Golden Delicious' apples (*Malus domestica* Borkh.) were subjected to either 0 C controlled-atm. (CA) storage or to a postharvest coating of 0.1% to 2.5% Nutri-Save (NS) a polysaccharide derived from shellfish plus air storage. NS-coated apples were greener and firmer and had higher titratable acidity (TA) and more shrivelled and injured fruit than the control after storage in air at 0 C for 5 to 6 months and ripening in air at 20 C for 7 days. Poststorage washing increased skin injury, and low rh during ripening increased shriveling of NS fruit. NS applications led to an accumulation of CO<sub>2</sub> and C<sub>2</sub>H<sub>4</sub> and a small reduction of O<sub>2</sub> in the fruit core cavities. The use of 1.5% O<sub>2</sub> + 1.5% CO<sub>2</sub> in the storage atm. was more effective than NS plus air storage in maintaining flesh firmness (FF) and TA without increasing fruit shrivel or skin injury. NS treatments maintained FF and a green skin in fruit ripened in air at 20 C for 2 or 4 wks following harvest, but some shrivel was evident by 4 wks. Better retention of skin greenness was the only benefit derived from a poststorage NS treatment of CA-stored fruit during the shelf-life test. AS

2026

Kramer (GF), Wang (CY) and Conway (WS). **Inhibition of softening of polyamine application in 'Golden Delicious' and 'McIntosh' apples.** *Journal of the American Society for Horticultural Science* 116(5): 1991; 813-817

2027

Andrich (G), Fiorentini (R), Tuci (A), Zinnal (A) and Sommovigo (G). **A tentative model to describe the respiration of stored apples.** *Journal of the American Society for Horticultural Science* 116(3): 1991; 478-481

Using mathematical equations that describe the O<sub>2</sub> mass-transfer and the enzymatic oxidation of the organic substrates of apples (*Malus domestica* Borkh.), a kinetic model to correlate fruit respiration rate with environmental O<sub>2</sub> partial pressure (PO<sub>2</sub>) was developed. The kinetic detn. were carried out at room temp. using apples stored at 3 to 4 C for 11 to 19 wks. Results show that: 1) the calculated value of the Michaelis-Menten constant related to the enzymatic oxidation of the respiratory substrate ( $K_m = 2.1 \text{ plus or minus } 0.5 \cdot 10^{-5} \text{ mol.kg}^{-1}$ ) is close to that reported in the literature for cytochrome-c oxidase; 2) the located range of PO<sub>2</sub> levels where O<sub>2</sub> becomes the limiting factor in the respiration process (near 2.6 kPa at T = 20.5 plus or minus 1 C)



is close to those usually used on a commercial scale for controlled atm. storage. AS

## Avocados

2028

Meir (S), Philosoph-Hadas (S), Zauberman (G), Fuchs (Y), Akerman (M), Aharoni (N). **Increased formation of fluorescent lipid-peroxidation products in avocado peels precedes other signs of ripening.** *Journal of the American Society for Horticultural Science* 116(5); 1991; 823-826

Fluorescent products (lipofuscin-like compounds) of lipid peroxidation, which accumulate with age, were extracted from 'Fuerte' avocado (*Persea americana* Mill.) peels during ripening. Fractionation and analysis of these fluorescent compounds (FCs) was carried out by an improved method, based on separation of FCs from chlorophyll by Sep-Pak silica cartridges. A sharp rise in FCs content was found 2 days after harvest in avocado fruits stored at 22 C, and ethylene enhanced this rise 3-fold on the 4th day. The accumulation of FCs preceded by at least 3 days the onset of climacteric ethylene and respiration and by 2 days the decrease in fruit firmness. Moreover, a 6-fold increase in the FCs concn. occurred during 1 to 2 wks of storage at 5 C, but the avocado fruits did not show any other detectable signs of ripening. These results suggest that lipid peroxidation may be regarded as one of the earliest detectable processes occurring during fruit ripening. Thus, an increase of FCs in peel may be employed as a horticultural characteristic for estimating initiation of ripening in avocado fruit. AS

## Citrus

2029

Chou (TC), Pintauro (N) and Kokini (JL). **Conformation of citrus pectin using small amplitude oscillatory rheometry.** *Journal of Food Science* 56(5); 1991; 1365-1368, 1371

Conformation of citrus pectin (uronide 93%, degree of esterification, 61.5%) sol. was studied and experimental reduced storage  $[G']_R$  and loss moduli  $[G'']_R$  were compared with random coil theories of Rouse and Zimm and the rodlike theory of Marvin and McKinney. The theory of Zimm best approximated experimental data. This suggested that components of the pectin molecule internally interact to form a compact random coil with conformation controlled by hydrodynamic interactions. AS

## Figs

2030

Pawar (SG), Kulkarni (DN), Shere (DM), Kulkarni (KD) and Patil (VK). **Effect of pretreatments on chemical composition and drying rates of solar dried figs.** *Indian Food Packer* 46(1); 1992; 39-44

Figs (*Ficus carica* L) var. Poona and Daulatabad were dried using solar drier after pretreating with (i) sulphitation, (ii) blanching + sulphitation, (iii) blanching, and (iv) lye treatment. Protein content was not altered by pretreatments. Total and reducing sugars were higher in sulphited Poona fig than in Daulatabad fig. Crude fiber was higher in Daulatabad. The drying rates at the initial stages were similar in all the treatments but at the final stages control and (ii) showed rapid drying rate in the Poona var. In the Daulatabad var., at the final stages of drying, control and (iii) showed faster drying rates. GS

## Grapefruits

2031

McDonald (RE), Miller (WR), McCollum (TG) and Eldon Brown (G). **Thiabendazole and imazalil applied at 53 C reduce chilling injury and decay of grapefruit.** *Hortscience* 26(4); 1991; 397-399

The fungicides thiabendazole (TBZ) or imazalil were applied at  $1 \text{ g.l}^{-1}$  at 24 or 53 C to 'Marsh' and 'Redblush' grapefruit (*Citrus paradisi* Macf.) to reduce fruit susceptibility to chilling injury (CI) and decay. Generally, there was more CI and decay on 'Marsh' grapefruit than on 'Redblush'. Severity of CI was lower in grapefruit that had been dipped at 53 C than at 24 C. Fruit dipped in fungicides had less CI than fruit dipped in water alone. Imazalil was more effective in reducing CI than TBZ. Fungicides reduced decay at both temp., and imazalil was better than TBZ. AS

## Grapes

2032

Sharma (PC), Joshi (VK) and Sharma (KD). **Preliminary studies on juice processing suitability of grapes grown under dry temperate conditions.** *Indian Food Packer* 46(1); 1992; 51-54

Juice processing suitability of three var. of grapes, (i) Katta, (ii) Chholtu Red, and (iii) Chholtu White grown under dry temperate conditions were evaluated. A suitable grape juice extraction technique was also standardised. Hot pressing gave better extraction of constituents like tannins, acids, total soluble solids and increased relative viscosity, compared to cold pressing. Clarification with pectolytic enzyme resulted in heavy settling of



sediments comprising fine pulp, tartarates and pectic substances at the bottom thereby improving the colour, body and sugar/acid/astringency blend of the juice. Extraction and clarification methods also affected the sensory qualities of the juices. Chholtu Red and Chholtu White juices were too acidic, astringent and unacceptable while Katta grape juice was superior to others, in colour, body and sugar/acid blend. GS

2033

Prakash (GS), Chadha (KL) and Reddy (BMC). **Changes in the physical and chemical composition of Kniffin trained grapes during ripening.** *Indian Journal of Horticulture* 48(4); 1991: 282-285

Changes in the berry wt., vol. and chemical constituents were determined in Thompson Seedless, Gulabi, Black Champa and Bangalore Blue grapes, trained to 6-arm Kniffin system at weekly interval from Veraison till maturity (January 28 to March 2). Berry wt. and size increased from Veraison till harvest in all cv. except in Bangalore Blue grape. Total soluble solids and its ratio with acids increased steadily from Veraison till maturity. The acid content in the berries decreased steadily after veraison until maturity. SRA

#### Mangoes

2034

Miller (WR), McDonald (RE) and Sharp (JL). **Quality changes during storage and ripening of "Tommy Atkins" mangoes treated with heated forced air.** *Hortscience* 26(4); 1991: 395-397

Freshly harvested mangoes (*Mangifera indica* L.) treated with forced air at 51.5 C for 125 min then stored for 1, 2, or 3 wks at 12 C, followed by 21 C until soft-ripe, were compared with nontreated fruit for quality changes. Treated fruit lost 1.0% more fresh wt. than nontreated fruit and developed trace amounts of peel pitting. Total soluble solids concn. for treated and nontreated fruit were similar (approx. 13%), as was peel colour at the soft-ripe stage. Treated fruit generally reached the soft-ripe stage approx. 1 day earlier than nontreated fruit regardless of storage duration and had a lower incidence and severity of stem-end rot and anthracnose. The trace of pitting on treated fruit likely will not influence consumer acceptance. AS

#### Peaches

2035

Nanos (GD) and Mitchell (FG). **Carbon dioxide injury and flesh softening following**

**high-temperature conditioning in peaches.** *Hortscience* 26(5); 1991: 562-563

High-temp. controlled-atm. (high CO<sub>2</sub>/low O<sub>2</sub>) conditioning was investigated as a possible treatment to delay the incidence of internal breakdown of peaches and nectarines (*Prunus persica* L. Batsch) during subsequent cold storage. Maintaining an atm. of 5% to 15% CO<sub>2</sub> added to air or to 1% to 5% O<sub>2</sub> while conditioning peaches for 2 days at 20 C partially prevented fruit ripening (compared to fruit conditioned in air), as measured by flesh softening and loss of green pigment, while no off-flavours were detected. Conditioning of peaches at 20 C for 4 days in air or in air + 20% CO<sub>2</sub> was detrimental to fruit quality, as indicated by flesh softening or detection of off-flavours. AS

#### Plums

2036

Sharma (KR) and Thakur (NS). **Evaluation of wooden boxes fabricated from lesser valued farm tree species for packaging and transportation of plums.** *Journal of Food Science and Technology (India)* 29(4); 1992: 235-236

'Santa Rosa' plums packed in the packages fabricated from lesser valued farm tree species were transported by truck to the market covering a distance of about 400 km. All the packing cases remained intact except cracking of one side strip in each single box fabricated from the woods of *Albizia chinensis*, *Bombax celba* and *Ficus roxburghii*. The quality of fruits was found to be acceptable with respect to loss in net and gross wts., fruit damage, fruit firmness and total soluble solids. AS

#### Strawberries

2037

Paakkonen (K) and Mattila (M). **Processing, packaging and storage effects on quality of freeze-dried strawberries.** *Journal of Food Science* 56(5); 1991: 1388-1392

The surface temp. during freeze drying (from 20 to 60 C) affected the hygroscopicity of the strawberries, particularly below 0.40 a<sub>w</sub>. Crushing before drying and low processing temp. improved sensory quality of dried strawberries and vacuum packaging improved storage stability. SRA

2038

Lesschaeve (I), Langlois (D) and Etievant (P). **Volatile compounds in strawberry jam: Influence of cooking on volatiles.** *Journal of Food Science* 56(5); 1991: 1393-1398



Volatile compounds in strawberry (*Fragaria ananassa* Duch) jam were only slightly affected by addition of sugar, but were closely related to design of the cooker and to the pressure used. Different types of behaviour were observed for the specific aromatic components. Cooker design largely influenced flavour losses at low pressures. Condensation of vapour during cooking and incorporation of the condensate in pectin sol. could result in a more flavourful product. SRA

## CONFECTIONERY, STARCH AND SUGAR

### Confectionery

#### Chocolates

2039

Joshi (NS) and Sharma (RS). **Milk chocolate.** *Indian Dairyman* 44(5): 1992: 225-230

The difference between plain chocolate and milk chocolate lies in the presence of milk solids and their sugar content. Cocoa nibs, cocoa mass, cocoa press cake, sugar, additives like emulsifiers, preservatives, flavourings and antioxidants all go into the making of milk chocolate. Use of cocoa butter is optional. Chocolate liquor is obtained by cleaning and grading, roasting, husking and winnowing and grinding of cocoa beans. This is mixed with powdered sugar, milk solids and cocoa butter. The chocolate mass is subjected to conching, storage, tempering, enrobing, moulding and packaging. GS

#### Sugar

2040

Rahmen (MH), Pal (SK) and Alem (F). **Effect of nitrogen, phosphorus, potassium, sulphur, zinc and manganese nutrients on yield and sucrose content of sugarcane (*Saccharum officinarum*) in flood-plain soils of Bangladesh.** *Indian Journal of Agricultural Sciences* 62(7): 1992: 450-455

The field exp. was carried out in the cropping season of 1986-87 to study the response of sugarcane to N, P, K, S, Zn and Mn fertilizers in sandy loam soils of sara series. The soil showed deficiency in N, P, S and Mn. It was found that the germination %, the number of tillers, millable canes, and yields of cane and sugar increased with the application of N, P, K, S, Zn and Mn at 150, 52, 83, 30, 8 and 5 kg/ha, respectively. Individual application of N, P, K, S and Zn increased the cane yield by 33.0, 34.4, 16.2, 16.2 and 8.7% respectively than the control. The highest

net benefit was obtained with application of N, P, K, S, Mn and Zn at the rate of 150, 52, 83, 30, 5 and 8 kg/ha respectively. KAR

2041

Patel (HS), Mehta (NJ), Patel (MP) and Vashi (RD). **Effects of farm yard manures, castor cake and press mud cake on yield and quality of sugarcane.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 93-102: 1991

The exp. conducted during the 1982-83 to 1984-85 indicated that farm yard manure (FYM) at 25 t/ha or castor cake at 625 kg/ha with 250 N, 125 P and 125 K/ha increased the cane yield, sucrose % juice and net return/ha. The treatments affected the sucrose % juice. Castor cake at 625 Kg/ha with the fertilizer gave higher sucrose % (17.84) and with max. commercial cane sugar % (12.20). This also gave max. additional net return of Rs.4251/ha. GS

2042

Banwari Lal. **Effect of nitrogen levels on growth, yield and quality of sugarcane under water-limited environment.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 109-114: 1991

The field exp. conducted in 1985-86 and 1986-87 involved growing sugarcane under 4 moisture regimes (unirrigated, 25% ASM, 50% ASM and 75% ASM) with application of 75, 150 and 225 kg N/ha. Sugarcane yield was significantly increased at 75% ASM regime by 115.68%, 89.00% and 41.50%, in 1985-86 and 128.20%, 89.01% and 40.19% in 1986-87 over unirrigated, 25% ASM and 50% ASM, respectively. The yield of sugarcane was also increased by 38.03%, 24.33% and 9.57% in 1985-86 and 64.42%, 40.61% and 16.79% in 1986-87 over control, with 75, 150 and 225 kg N/ha, respectively. The quality of sugarcane did not change due to different soil moisture regimes or different N levels. GS

2043

Onkar Singh and Kanwar (RS). **Effect of crop geometry on physiological activity and juice quality of sugarcane.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 141-147: 1991

Sugarcane crop was grown with a geometry of E. to W. and N. to S. directions of rows with 60, 90 and 120 cm spacing between rows. The physiological activity and sugarcane juice quality of cv. CoJ 64 were better under 120 cm row spacing than 60 cm spacing. The crop geometry of E. to W. row direction with 120 cm row spacing is considered to be better



than N. to S. directions of rows and 60 and 90 cm spacings between rows. The sucrose, purity, CCS, electric conductivity and pH were slightly higher while reducing sugars, fibre content, gums and flavonoids of juice were lower under E. to W. than N. to S. direction of rows. GS

2044

Bhojaraj (SK). **Development of poly baffle entrainment catcher on effective device to prevent entrainment in sugar factory.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 1-14; 1991

Prevention of entrainment and arresting of entrained sugar liquors from the evaporator bodies and vacuum pans of the sugar factories has assumed greater importance. The poly baffle entrainment catcher, which is the most advanced technique comprises of multi baffle plate made of stainless steel 304 quality as a corrugated sheet having a vortex at the top and steep angle between two layers. The vapour travelling through the arrestor strikes on the corrugated surface of the baffles which changes its direction resulting in the falling of droplets of the entrained sugar liquor. This solves the problem of entrainment and the allied losses in sugar recovery. GS

2045

Jain (PF), Ahire (CN), Joshi (JT) and Baranth (AM). **Power reduction in condensing system.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 15-28; 1991

In multijet spray condenser, cold water is provided to both spray and jet nozzles. In the modified system, instead of cold water, hot water was provided through the condenser outlet. By this, quality of water to be handled in spray pond reduced and one spray pump could be stopped, which resulted in power reduction also. Expenditure involved in the modification is Rs. 100,000 and the saving in the power consumption is Rs. 117,000. GS

2046

Dixit (AN). **Optimum use of exhaust steam desuperheaters in sugar industry.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 29-38; 1991

Optimum use of desuperheaters to reduce superheated process steam temp. in sugar industries has been considered. Fixing of capillary type good quality temp. gauge in incoming and outgoing steam in desuperheater, installing two recorder type flow meters, one in superheated steam line and other in water line to know actual flow of

both fluids are suggested. The ratio of peripheral area of desuperheater to cross sectional area of incoming pipe should be 1.25 only. Pressure maintained at Header should be 150 psig. These modifications will yield desired results in the sugar industry. GS

2047

Phatak (RN) and Malshe (VC). **Condensate water treatment in sugar industries.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 39-45; 1991

Lowering of boiler water pH, when mixed with condensate water (for economy) is a problem faced by sugar industries. All condensate samples had pH value near neutral or alkaline. But after boiling the condensate water the contaminants (sugar or its derivatives) decompose into acidic components leading to low pH. On boiling with or without caustic, variation was observed in pH value. But boiling with 50 p.p.m. caustic showed very little effect. Addition of oxidizing agents at predetermined rate (0-20 p.p.m.); scavenger resin treatment; and desorption of organics to regenerate the resin maintained the desired pH of water. GS

2048

Kapur (PPatilVL), Gautam (GK), Vasudeva (TR), Agarwal (AK), Sai Kumar (KS), Bajaj (VD). **On-line estimation of crystallisation parameters and automatic feed control in vacuum pans.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 17-27; 1991

The salient features of the microprocessor based pan boiling instrumentation system (MIPMOS) developed by the Central Electronics Engineering Research Institute (CEERI), Pilani, specifically for low grade vacuum pans of the sugar industries, have been described. MIPMOS has helped in higher productivity, energy conservation and minimisation of losses with crystal quality improvement. The hardware and software configurations; specifications of the system, and the data collected during the field operation of the system are given. GS

2049

Bhupinder (K) and Sharma (KP). **Techno-economic development of agro-industrial based products. I. "Ready-to-serve" sugarcane juice (pure).** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 39-50; 1991

Ready-to-serve semi-processed (pasteurized and preservative added) and processed (preservative



added and sterilized) sugarcane juice, was analysed using regression techniques, during 120 days storage at 10 C, room temp. and 40 C, to find the rate of change of sugar loss, acidity development, colour and acceptability index. The rate of sucrose loss per day in stored juice was 0.042% for processed and 0.327% for semi-processed juice stored at room temp. The loss of quality of processed juice was slower than semi-processed juice during 120 days of storage at 10 C and at room temp. GS

2050

Laxmi Narayan, Rao (KSN) and Srinivas (K). **Microprocessor based control system for mixed juice flow.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 51-58; 1991

Mixed juice flow in sugar mills affects the primary heating, juice clarification, secondary heating and further processing stages. Hence, it is important in energy conversation and to achieve process efficiency. The salient features of microprocessor based flow control system using magnetic flow meter for controlling the mixed juice flow was developed and tried in sugar factory. GS

2051

Dubey (HK), Dubey (Y) and Sharma (GC). **Effect of different growth stimulating supplements on molasses fermentation by distillers yeast.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 59-63; 1991

In molasses fermentation by yeast, growth stimulating supplements like progesterone, estrogen and oxytocin stimulated the growth, sugar utilization and the rate of alcohol production and yield of alcohol. Fermentation efficiency was 93.5% for progesterone, 94.7% for estrogen, and 94.92% for oxytocin at 30 C and at 2 p.p.m. level. GS

2052

Srinivasan (S), Gunasekaran (N), Gurugovind (J) and Chandrasekar (N). **Performance of a falling film evaporator as first effect.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 1-16; 1991

Falling film vessel (FFV) when compared with Roberts type FFV, showed better steam economy and less of scaling with better efficiency. Other advantages include no expenditure on energy for upward movement of the juice, negligible retention time and avoidance of sugar loss; no boiling point elevation; no loss of temp. difference; no juice stagnation on the heating surface; provision of 75%

recirculation of the juice; and no need to increase the exhaust pressure. GS

2053

Murthy (TSN). **Theory and practice of cane juice clarification in sulphitation sugar factories.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 17-23; 1991

For juice clarification in sulphitation plants important aspects to be considered include quality and quantity of lime, temp. control, sulphur dioxide, pH, Brix and use of some auxillary chemicals in getting better clarified juice or making the mud to settle faster. These are described. GS

2054

Kaliyamurthy (S) and Ramasamy (P). **Simultaneous achievement of good colour sugar and lowest final molasses purity - a novel idea.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 25-36; 1991

A new method to get lowest final molasses purity and good colour sugar is suggested. 'B' massecuite is cured in a continuous centrifugal machine without adding any water, to maintain the purity at 36. The 'B' fore sugar will be made into 'B' magma by the addition of 'B' light and this 'B' magma is cured by another continuous centrifugal machine with sufficient water washing. The additional requirement of machineries for this process is a centrifugal machine, pugmill, Mingler, magma pump, 'B' light pump and 'B' light tank at the pan floor. By the double curing of 'B' massecuite, 36 purity 'B' heavy can be had; with 'C' massecuite of 45 purity; 22, 23 purity of final molasses and good 'A' quality sugar. GS

2055

Verma (VK) and Tayagi (ID). **Selection of optimum pH of juice in juice sulphiter.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 37-46; 1991

The selection of optimum pH of sugarcane juice in juice sulphiter is done by providing an additional connection for lime in juice sulphiter by maintaining the initial pH of juice at 8.6 and by neutralising it to 7.1 pH by SO<sub>2</sub>. This helped to reduced CaO contents in clear juice, less destruction of reducing sugar and better rise in purity from mixed juice to clear juice. It also improved the sugar colour and reduced sugar loss in molasses. GS



2056

Chauhan (SS). **Automation on juice sulphitation by controlling of SO<sub>2</sub> gas generation, a new approach.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 47-58; 1991

In juice clarification, manual control of pH gave variable results of clarification efficiency and chemical requirements. An automatic control system on clarification process has been developed. This involved the control of the SO<sub>2</sub> gas generation and lime dose. Auto control of SO<sub>2</sub> generation has been done by controlling the air supply to the burner with minor modification in the existing system. This has resulted in reduced sulphur and lime consumption, and increased the clarification efficiency and reduced the cleaning period. GS

2057

Shukla (BK). **Review of basic absolute sugar recovery and sugar recovery performance.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 59-65; 1991

The basic absolute sugar recovery (BASR) is used by the author to arrive at basic sugar recovery based on certain basic assumptions. The term represents 100% mills extraction and the basic sugar recovery is calculated on the basis of absolute juice % cane and its calculated value of Pol % (available boiling house recovery) and Brix % absolute juice based on certain assumption and a virtual final molasses purity 28.57. Sugar recovery is divided by BASR. Thus at the absolute juice purity of 85, Pol in cane is equal to 12.5% and fibre % in cane is equal to 12.5, the max. BASR is 11.625. This will give (BASR) the max. of 93% basic overall recovery. GS

2058

Agarwal (JKP) and Agarwal (S). **Nature of losses in a sugar factory.** *Proceedings of the 53rd Annual Convention of the Sugar Technologist's Association of India* 67-72; 1991

Describes known, unknown and undetermined losses such as mechanical losses due to entertainment, leakage and spillage, incorrect weighing or measurement; chemical losses due to inversion of sucrose into reducing sugars which may be due to microbiological growth in mill beds, juice pipes, centrifugal flooring; sugar loss in evaporator from syrup to sugar stage; loss in bagasse; loss in final molasses and loss in press cake. Methods for calculating the above losses are indicated. GS

2059

Kanda (K), Yasuda (Y) and Tochikubo (K). **Germination response of *Bacillus subtilis* PC1219 spores to caramelized sugar and L-asparagine.** *Journal of Food Science* 56(5); 1991: 1399-1403

In presence of L-asparagine, effective substances in caramelized sugar were primarily glucose and fructose; isomerization of glucose/fructose was observed after autoclaving (120 C, 20 min, pH 7.2). Glucose and fructose were 63 to 24 in caramelized glucose and 12 to 44 in caramelized fructose. Apparent dissociation constants were: glucose  $2.2 \times 10^{-4}$ , fructose  $1.3 \times 10^{-4}$ , L-asparagine  $2.2 \times 10^{-4}$ . Hydroxymethylfurfural and maltol were not effective. Aging and heat activation of spores contributed to germination, especially for response to L-asparagine and/or fructose, but not glucose. The initiation mechanism was distinguishable from the L-alanine system in requirement for aging, heat activation of spores and response to inhibitor. L-Asparagine could be partially replaced by NH<sub>4</sub><sup>+</sup>, but not by L-glutamine. AS

## BAKERY PRODUCTS

2060

Abbott (J), O'Palka (J) and McGuire (CF). **Dried distillers' grains with solubles: Particle size effects on volume and acceptability of baked products.** *Journal of Food Science* 56(5); 1991: 1323-1326

Particle size distribution of sour mash dried distillers' grains solubles (DDGS) varied among producers but did not influence final vol. of quick muffins or yeast rolls. Vol. of muffins or rolls was not affected by sieve fractions, particle size distribution, grinding effects or DDGS source. Ground DDGS had little effect on baked product colour. Quick breads and yeast breads baked with DDGS produced darker colours than standard bread without DDGS. Consumer acceptability of muffins or rolls was not altered by DDGS additions (P greater than or equal to 0.05). SRA

## Bread

2061

Vatsala (CN) and Haridas Rao (P). **Studies on fruit bread.** *Journal of Food Science and Technology (India)* 29(4); 1992: 213-217

Studies on fruit bread indicated that min. levels of 15.5 and 25% sugar, fat and fruit respectively were required to obtain bread with desired taste. The bread thus made had slightly undesirable hard



texture which was overcome by increasing the fermentation and proofing times from 120 to 150 and 55 to 70 min respectively. Fruit bread dough required about 6% less water as compared to that required for a normal bread dough. Fermentation rate was reduced as indicated by the Expansograph characteristics showing longer time to reach the peak (135 min) compared to normal bread dough (65 min). Fruit bread could be stored at ambient conditions of 25 - 28 C and rh of 60 - 65% after packing in polypropylene pouches (100 gauge) for a longer period of 8 days without mould growth and with min. of textural changes in contrast to that for 3 days for normal bread. AS

## Cakes

2062

Bell (DA) and Steinke (LW). **Evaluating structure and texture effects of methylcellulose gums in microwave-baked cakes.** *Cereal Foods World* 36(11); 1991; 941-944

Methylcellulose gums significantly improve microwave cake height, texture and moisture retention. Special testing procedures used to evaluate their effects provided statistically significant relationships between the measured responses and the independent variables, gum concn. (0 - 4% of flour wt.) and moisture added in excess of control (0 - 30% of flour wt.). BV

## MILK AND DAIRY PRODUCTS

2063

Donnelly (WJ). **Applications of biotechnology and separation technology in dairy processing.** *Journal of the Society of Dairy Technology* 44(3); 1991; 67-72

The impact of biotechnology on dairy processing within the next decade are reviewed. Microbial processes, lactose fermentation, new enzyme applications (milk protein hydrolysates, special peptides, milk fat modification), accelerated cheese ripening, milk protein fractionation and antimicrobial proteins are the topics covered in this review. BV

2064

Aguilera (JM) and Kinsella (JE). **Compression strength of dairy gels and microstructural interpretation.** *Journal of Food Science* 56(5); 1991; 1224-1228

Contour plots were developed for the compression stress (at 20% deformation) of single-component,

mixed and filled protein gels. Samples were made by heating and acidification from skim milk powder, SMP (0-20% TS), whey protein isolate, WPI (0-10% TS), and recombined cream, within pH 3.6 - 3.9, 4.6 - 4.8 and 5.1 - 5.3. At higher pH, WPI gels were stronger than SMP gels. WPI had a reinforcing effect on SMP gels, while small additions of SMP to WPI gels resulted in weaker mixed gels. Filled gels containing cream had higher compression strengths than mixed gels. Micrographs showed linking of casein chains by WPI strands in mixed gels and compatibility of fat globules with casein micelles in the protein network of filled gels. AS

## Milk

2065

Nagarajan (L), Selvakumaran (R), Kumar (VU) and Umesh Kumar (S). **Deterioration in milk quality due to a psychrotrophic *Bacillus cereus* during storage at low temperatures.** *Chemie Mikrobiologie Technologie der Lebensmittel* 12(6); 1990; 168-170

Psychrotrophic *Bacillus cereus* survived in milk during pasteurization and grew at 4 C storage and increased proteolytic activity. Results indicate deterioration of milk quality due to *B. cereus*. BV

2066

Chavan (KD) and Kulkarni (MB). **Studies on processing properties of milk obtained from crossbred cows.** *Journal of Food Science and Technology (India)* 29(4); 1992; 245-246

The data indicated that the processing qualities of the milk are influenced by the genetic group of cows in terms of rennet and heat clotting times as well as curd tension. Microbial rennet is advantageous for quick coagulation while the milk from HF crossbred cows is more suited for formulating infant foods. AS

2067

Kanno (C), Shimomura (Y) and Takano (E). **Physicochemical properties of milk fat emulsions stabilized with bovine milk fat globule membrane.** *Journal of Food Science* 56(5); 1991; 1219-1223

Milk fat globules (MFG) were reconstituted with milk fat globule membrane (MFGM) and milk fat (MF). Viscosity of the reconstituted MFG was highest at pH 5.0 and 4 min emulsifying, and rose with an increase of MFGM between 40 - 80 mg/g fat. Adsorbed protein/unit fat increased at acid pH with increase of MFGM. The composition of proteins adsorbed on the surface of MFG was not influenced by factors of reconstitution. The size and specific surface area of globules were influenced by



emulsifying time, MFGM and MF concn., and pH. The size range of MFG prepared by standard method was 0.9 - 17  $\mu\text{m}$  in diam. Median diam. was 5  $\mu\text{m}$  and specific surface area was 15,600  $\text{cm}^2/\text{cm}^3$  of emulsion. AS

## Milk products

2068

Samona (A) and Robinson (RK). **Enumeration of bifidobacteria in dairy products.** *Journal of the Society of Dairy Technology* 44(3): 1991: 64-66

A range of selective and non-selective media for enumeration of bifidobacteria in dairy products showed that modified Rogosa agar method is the best for recovery from yoghurt-like products. BV

2069

Fu (B), Taoukis (PS) and Labuza (TP). **Predictive microbiology for monitoring spoilage of dairy products with time-temperature integrators.** *Journal of Food Science* 56(5): 1991: 1209-1215

Arrhenius and square root equations were used to model the growth of *Pseudomonas fragi*. Significant negative history effects for growth rate and positive effects on lag phase of *P. fragi* were found under nonisothermal conditions. A correlation scheme of *P. fragi* growth was developed with the time/temp. integrators (TTI) response. Both the Arrhenius and square root model fit the lag phase and the growth rate data of the microbe at constant temp. very well. Both would be needed for shelf-life prediction. The application of TTI's for dairy products is feasible despite the history effects. SRA

2070

Desai (NB), Desai (HK) and Thakar (PN). **Response surface methodology vis-a-vis optimization of sensory quality of dairy products.** *Indian Dairyman* 44(5): 1992: 237-240

The review covers Response Surface Methodology (RSM) as a technique for improving probability of success of products in the market. The 4 steps involved in RSM, considerations while making use of RSM and application of RSM in arriving at optimal sensory quality and in setting sensory standards have also been covered. 3 references. GS

## Butter

2071

Schooner (F), Simard (RE) and Pandian (S). **Colorimetric assay for free fatty acids in butter**

**using flow-injection and immobilized enzymes.** *Journal of Food Science* 56(5): 1991: 1229-1232

Butyrate kinase (BK) was extracted from *Escherichia coli* DH5 pJc7\*, purified, immobilized on porous glass beads and packed in a glass tube as a biosensor coupled with a flow-injection analysis system (FIA). The limits of detection were: FIA (25 p.p.m.) and batch method (BM, 5 p.p.m.). BK is highly specific for butyric, propionic and valeric acids. Regression coeff. values were: FIA  $r = 0.992$  and free enzyme  $r = 0.987$ . The FIA and BM were tested on butter samples with high correlation ( $r = 1.000$ ) between the methods. Correlation coeff. were:  $r = 0.994$  and  $r = 0.988$  between acid degree value and FIA/BM. AS

2072

Garcia (HS), Amundson (CH) and Hill (CGJr). **Partial characterization of the action of an *A. niger* lipase on butteroil emulsions.** *Journal of Food Science* 56(5): 1991: 1233-1237

A commercially available lipase (*Aspergillus niger*, APF-12) was employed to hydrolyze an emulsified butteroil substrate. HPLC analyses of the reaction products indicated that short chain fatty acid residues (e.g., butyric acid) were preferably hydrolyzed at pH 5 whereas the overall rate of hydrolysis for all major fatty acids showed a pH optimum between 6 and 7. This indicated the possibility of directing the selectivity of lipolysis in butteroil to enhance production of short-chain fatty acids associated with flavour development. The optimal temp. was ca. 35 C. Different cations also affected rate of hydrolysis. AS

## Cheese

2073

El-Abbassy (MZ), Aly (ME) and Taha (SN). **Effect of lecithin addition on the yield and quality of soft cheeses.** *Die Nahrung* 35(6): 1991: 633-640

The effect of lecithin added at levels of 0.025, 0.050 and 0.075% to cheese milk used in manufacture of Domiati and Kareish cheese on yield, wt. losses and quality of cheese was studied. Addition of lecithin increased cheese yield and decreased wt. losses during pickling. The bacterial content of all cheeses treated with lecithin was higher than that of control cheese when fresh and during pickling. Cheese made with added lecithin showed higher moisture, salt, fat and acidity than control cheese. The total N percentage was almost the same in all treatments. The levels of both soluble N and total volatile fatty acids in cheese containing lecithin were higher than in the control cheese. No marked differences were evaluated for flavour, body and texture of fresh



cheese made from milk with or without lecithin, while during pickling the organoleptic properties of cheese containing lecithin were improved. AS

2074

Ha (JK) and Lindsay (RC). **Volatile branched-chain fatty acids and phenolic compounds in aged Italian cheese flavours.** *Journal of Food Science* 56(5): 1991: 1241-1247, 1250

Analysis of cheeses by capillary GC revealed adequate concn. of certain branched-chain fatty acids (BCFAs) and phenols present to contribute distinctive flavours. Butyric acid-like flavours in cow's milk Romano cheese were modified by 2-methyl-butanoic and 2-ethylbutanoic acids which provided sweet, fruity notes. 4-Ethyl-octanoic acid provided a characterizing goaty note to Romano cheese made from mixed goats' and cows milk. 4-methyl-octanoic and 4-ethyl-octanoic acids along with *p*-cresol, *m*-cresol, and 3,4-Di-methylphenol appeared responsible for sheepy notes in sheep milk Romano cheese. Phenol and cresols (o,m,p) strongly contributed phenolic and medicinal flavour notes to smoked Provolone cheese. Low concn. of BCFAs and phenols appeared to provide desirable background flavours to Parmesan cheeses. AS

2075

Chandan (RC). **Cheeses made by direct acidification.** *Indian Dairyman* 44(4): 1992: 180-188

Latin American White (LAW) cheeses are made from acidified whole milk, skim-milk, cream or mixture thereof. Ripening, shelf-life, composition, nutrient, density and uses of LAW cheeses are described. *Chhana* which is also a common acidified cheese is described with its production techniques, and its utilization in preparing *sandesh* and *rasogolla*. GS

### Cheddar cheese

2076

Trepanier (G), Simard (RE) and Lee (BH). **Lactic acid bacteria relation to accelerated maturation of Cheddar cheese.** *Journal of Food Science* 56(5): 1991: 1238-1240, 1254

2077

Tewari (BD), Vipin Kumar and Singh (S). **Spray drying process for accelerated ripened Cheddar cheese.** *Indian Dairyman* 44(5): 1992: 244-248

Method for manufacture of dried cheese from Cheddar cheese and accelerated ripened curd slurry has been described. Preparation of slurry for spray drying consists of selection of cheese, blending,

addition of emulsifying salts, water and other additives followed by heat processing and homogenization. The slurry is spray dried, cooled, packed and stored as powder. The cheese powder from Cheddar cheese has got bulk density 0.46 g/ml, dispersibility 34.20 g, flowability 37°, sinkability after 6 min in terms of % absorbance 1.0 and the corresponding values for powder made from accelerated ripened curd are 0.46 g/ml, 29.34 g, 40° and 0.85 respectively. Slurry obtained from blend A (cow milk Cheddar cheese ripened with microbial rennet for 8, 4-6 and 2-3 months is blended in the ratio of 1:2:1) has higher pH, titratable acidity, soluble protein, but is lower in its salt and total volatile fatty acid than that from blend B (microbial rennet, cow milk Cheddar cheese (8 months old) and accelerated ripened curd blended in the ratio of 1:3). Reconstituted cheese spread could be prepared by mixing 2 parts of powder with 3 parts of water at temp. of 50 - 55 C and indirectly heating to 65 C with constant stirring. The product is allowed to set at 8-10 C for about 3-4 min before it is served. GS

### Kachhagolla

2078

Sen (DC). **Kachhagolla - a delicious milk sweetmeat of Bengal.** *Indian Dairyman* 44(5): 1992: 249-251

Preparation of *Kachhagolla* involves *chhana* prepared from cow's milk is kneaded and mixed with cane sugar at 30 to 35% by wt. of *chhana* in an iron karahi and cooked with controlled heating and stirring. It develops its characteristic sticky granular texture and aroma. In *kachhagolla*, moisture, fat, protein, sucrose, ash, titratable acidity, water soluble acid, free fatty acidity and free fat contents vary with the source of the market sample. Ordinarily the major constituents will have fat 6.14 - 9.54%, protein 10.32 - 11.26%, moisture 25.14 - 28.73% and sucrose content of 29.58 - 34.76%. *Kachhagolla* is packaged in folding paper board cartons and it has a limited shelf-life of 1 to 2 days in normal and between 3 and 5 days in refrigerated temp. GS

### Rasogolla

2079

Tambat (RV), Khorgade (AB), Changade (SP) and Kaloti (SV). **Effect of fat and maida levels on rasogolla preparation.** *Indian Dairyman* 44(4): 1992: 203-205

Cow milk was standardised to 3, 3.5, 4 and 4.5% fat levels and converted to *chhana* using 1.5% lactic acid sol. During *rasogolla* preparation, maida levels were 0, 2, 4, 6 and 8% on wt. basis of *chhana*.



Sensory score for colour and appearance, body and texture, flavour and taste and overall acceptability of *rasogolla* was evaluated on a 25 point scale. No remarkable changes in sensory score was observed due to various fat levels and were non significant. However, 2 and 4% maida levels recorded the highest score and were at par. Combined effect of fat and maida levels were non significant. *Chhana* yield, moisture content, fat losses in whey and coagulant sol. required for *chhana* preparation were increased with the increase of fat levels in milk. AS

## Wheys

2080

Colbert (LB) and Decker (EA). **Antioxidant activity of an ultrafiltration permeate from acid whey.** *Journal of Food Science* 56(5): 1991; 1248-1250

Results of the study indicate that both acid whey and acid whey ultrafiltration (UF) permeate could be used as "natural" antioxidants in foods. Acid whey inhibited up to 90% of Fe-catalyzed oxidation phosphatidylcholine liposomes. Lipid oxidation catalysts, including Fe, lipoxidase, photoactivated riboflavin and hydrogen peroxide-activated met Mb, were inhibited by acid whey and its UF permeate. The antioxidant activity of the UF permeate increased with increasing pH (5.4 - 7.0). Use of acid whey and its UF permeate as an antioxidant would increase utilization of a major by-product of the cheese industry and would decrease whey disposal problems. SRA

## Whey protein concentrates

2081

Kilara (A) and Mangino (ME). **Relationship of solubility of whey protein concentrates to thermal properties determined by differential scanning calorimetry.** *Journal of Food Science* 56(5): 1991; 1448-1449

Twelve samples of whey protein concentrates manufactured under defined conditions were analyzed for their composition. Samples were subjected to differential scanning calorimetry to determine the onset temp. and enthalpy of denaturation. Solubility of the samples at pH 7.0 was determined using three procedures. In each instance, solubility significantly correlated with both onset temp. and enthalpy of denaturation. Enthalpy of denaturation may serve as a predictor for solubility of proteins. AS

## Milk proteins

2082

Hung (SC) and Zayas (JF). **Emulsifying capacity and emulsion stability of milk proteins and corn germ protein flour.** *Journal of Food Science* 56(5): 1991; 1216-1218, 1223

Comparative studies of corn germ protein flour (CGPF), non-fat dry milk (NFDM), whey protein concentrates (WPC), and sodium caseinate (SC) showed that they were effective protein sources in the model system in terms of emulsifying capacity (EC) and emulsion stability (ES). EC and ES of all samples increased with increased concn. of protein. A max. EC at pH 8.0 was obtained for CGPF, WPC and SC while NFDM showed a max. EC at pH about 7.0. Minimal pH effect on ES was found. No effect of incubation temp. and incubation time on EC and ES of these samples was detected. SRA

2083

Strange (ED), Van Hekken (D) and Thompson (MP). **Qualitative and quantitative determination of caseins with reverse-phase and anion-exchange HPLC.** *Journal of Food Science* 56(5): 1991; 1415-1420

Whole native caseinate (WNC) and casein (CN) fractions from preparative DE-52 cellulose urea columns were chromatographed using C-8 reverse-phase (RP) and DEAE-type anion exchange (AEx) HPLC systems. With RP,  $\alpha_{s2}$ -CN and  $k$ -CN eluted first as several small peaks;  $\alpha_{s1}$ -CN eluted later as two peaks, followed by  $\beta$ -CN peaks. With AEx,  $k$ -CN eluted early as a group of peaks,  $\beta$ -CN eluted next, and  $\alpha_{s1}$ -CN and  $\alpha_{s2}$ -CN coeluted last. Standard curves were prepared for  $\alpha_{s1}$ -CN and  $\beta$ -CN using RP-HPLC and showed correlation coeff. of 0.99 and 0.98 respectively. The caseins in WNC, nonfat dry milk casein, commercial casein(ates) and caseins from milks of individual cows were determined. AS

## MEAT AND POULTRY

2084

Madsen (M), Milne (JAC) and Chambers (P). **Critical control points in the slaughter and dressing of farmed crocodiles.** *Journal of Food Science and Technology (India)* 29(4): 1992; 265-267

2085

Pommier (SA). **Characterization of the increase in paleness of milk-fed veal during refrigerated storage.** *Journal of Food Science* 56(5): 1991; 1455-1456

Ninety-five Holstein bull calves raised on milk replacer were used to study the development of meat



colour during storage. Luminous reflectivity (Y) of the pectoralis profundus muscle increased an average of 22 lux units after a 3-day aging period. Forty-five-minute colour measurements correlated with a 3-day colour measurements ( $r = 0.69$ ) and pre-slaughter hemoglobin ( $r = 0.53$ ), which also correlated with 3-day colour measurements ( $r = 0.65$ ). Meat colour at 45 min and preslaughter haemoglobin concn. contributed significantly to predict 3-day Y values ( $R^2 = 0.59$ ). However, this procedure was not accurate to predict colour of meat at 3 days after slaughter. AS

2086

Appu Rao (AG). **A stoichiometric analysis of bovine serum albumin-gossypol interactions: A fluorescence quenching study.** *Indian Journal of Biochemistry and Biophysics* 29(2); 1992: 179-182

The interaction of gossypol with bovine serum albumin, human serum albumin and n-bromosuccinimide-modified bovine serum albumin has been followed by fluorescence quenching measurements. The presence of a high affinity site (association constant  $K = 2.2 \times 10^6 \text{ M}^{-1}$ ) for gossypol on bovine serum albumin and human serum albumin is indicated. The stoichiometry of binding for the high affinity site was evaluated using Job's method of continuous variation, thereby suggesting the formation of 1:1 complex. Modification of the tryptophan residues on bovine serum albumin does not affect the binding of gossypol to either high or low affinity site of albumin. AS

## Meat

2087

Park (YW), Kouassi (MA) and Chin (KB). **Moisture, total fat and cholesterol and goat organ and muscle meat.** *Journal of Food Science* 56(5); 1991: 1191-1193

Moisture, total fat and cholesterol in liver, kidney, heart, *Longissimus dorsi* (LD) and *Biceps femoris* (BF) muscles of 2 goat breeds (Alpine and Nubian) were determined; and differences in lipids between these tissues as well as between breeds of goat were compared. Samples from goat kids fed with a complete diet containing 4 levels of Ca for 12 wk before slaughter were used for the study. Ranges of moisture (%) in liver was 68.4 - 82.9, kidney 75.8 - 85.0, heart 69.7 - 84.3, LD 74.8 - 83.2, and BF 76.0 - 84.7. The mean total fat % was liver 3.26, kidney 3.14, heart 4.32, LD 2.27 and BF 2.03 for pooled data of two breeds. The difference in cholesterol between liver and kidney was significant ( $P < 0.01$ ). The two skeletal muscles contained significantly ( $P < 0.01$  or  $0.05$ ) less cholesterol than all organ

tissues. Heart contained less cholesterol ( $P < 0.05$ ) than liver or kidney. Fat and cholesterol were influenced by dietary Ca, breed and tissue, while moisture was affected only by tissue. SRA

2088

Ha (JK) and Lindsay (RC). **Volatile alkylphenols and thiophenol in species-related characterizing flavours and red meats.** *Journal of Food Science* 56(5); 1991: 1197-1202

2089

Shahidi (F) and Pegg (RB). **Novel synthesis of cooked cured-meat pigment.** *Journal of Food Science* 56(5); 1991: 1205-1208

Cooked cured-meat pigment was synthesized from bovine blood and a nitrosating agent in aqueous sol. at elevated temp. Effects of reductant(s) and base in the reaction mixture were investigated on the yield and purity of the pigment. The best yield of pigment was 95% with purity  $> 98\%$  in most cases. Absorption characteristics of the pigment were identical to those of pigments extracted from a nitrite-cured ham or from hemin-nitric oxide synthesis. Colour imparted to comminuted pork by the pre-formed pigment, upon cooking, and its colour stability thereafter, were indistinguishable from that of nitrite-cured meat, yet depended on the myoglobin content. AS

2090

Shahidi (F), Pegg (RB) and Shamsuzzaman (K). **Colour and oxidative stability of nitrite-free cured meat after  $\gamma$ -irradiation.** *Journal of Food Science* 56(5); 1991: 1450-1452

The effects of 5 and 10 kGy irradiation on the colour and oxidative stability of meats treated with nitrite or a nitrite-free curing system were investigated. Irradiation had no detrimental effects on the colour or flavour of either cured samples. Polyphosphates had a beneficial effect on oxidative stability but had a slight detrimental effect on colour stability of irradiated samples. BV

2091

Palombo (R), Van Roon (PS), Prins (A) and Krol (B). **Comparative study on three methods for determining air content in meat batters.** *Journal of Food Science* 56(5); 1991: 1453-1454

Measurement of air content in meat batters is useful, as larger amounts of air negatively affect some properties of comminuted products. Three methods were compared. One method (Vemag) involved liberation of entrapped air from the batter by the combined effects of vacuum and mechanical



forces. The other two methods involved application of under- (Smrat) or over- (Handtmann) pressure and calculation of the air content using Boyle's ideal gas law. The Smrat showed the lowest reproducibility. However, being the most rapid and easy to use, it was the method of choice for routine quality and process control. The Vemag and more favourably the Handtmann were recommended for accurate and reproducible detn. AS

## Beef

2092

Kim Ha (J) and Lindsay (RC). **Volatile fatty acids in flavours of potatoes deep-fried in a beef blend.** *Journal of the American Oil Chemist's Society* 68(5): 1991: 294-298

The volatile free fatty acids were analyzed in commercial french-fried potatoes that had been deep-fried in beef tallow-hydrogenated vegetable oil shortening. The results showed that many of the volatile fatty acids present in beef tallow were transferred to the potatoes. Of the fatty acids derived from beef tallow, butanoic, 2-methylbutanoic, 3-methylbutanoic, heptanoic, 4-methyltanoic, and nonanoic acids were present in concn. above their respective thresholds, and should contribute to fallow-like flavours. Several unsaturated volatile fatty acids also were identified, and they should contribute to the general deep-fried potato flavour. AS

2093

Mitchell (GE), Giles (JE), Rogers (SA), Tan (LT), Naidoo (RJ), Ferguson (DM). **Tenderizing, ageing and thawing effects on sensory, chemical, and physical properties of beef steaks.** *Journal of Food Science* 56(5): 1991: 1125-1129

2094

Shackelford (SD), Koohmaraie (M), Whipple (G), Wheeler (TL), Miller (MF), Crouse (JD), Reagan (JO). **Predictors of beef tenderness: Development and verification.** *Journal of Food Science* 56(5): 1991: 1130-1135, 1140

2095

Payne (CA), Moody (WG), Langlois (BE), Means (WJ) and Aaron (DK). **Microbial characteristics of three formulations of precooked, vacuum-packaged restructured beef steaks.** *Journal of Food Science* 56(5): 1991: 1136-1140

2096

Papadopoulos (LS), Miller (RK), Acuff (GR), Lucia (LM), Vanderzant (C), Cross (HR). **Consumer and trained sensory comparisons of cooked beef top**

**rounds treated with sodium lactate.** *Journal of Food Science* 56(5): 1991: 1141-1146, 1153

2097

Stutz (HK), Silverman (GJ), Angelini (P) and Levin (RE). **Bacteria and volatile compounds associated with ground beef spoilage.** *Journal of Food Science* 56(5): 1991: 1147-1153

2098

Minerich (PL), Addis (PB), Epley (RJ) and Bingham (C). **Properties of wild rice/ground beef mixtures.** *Journal of Food Science* 56(5): 1991: 1154-1157

2099

Miller (AJ) and Menichillo (DA). **Blood fraction effects on the antibotulinal efficacy of nitrite in model beef sausages.** *Journal of Food Science* 56(5): 1991: 1158-1160, 1181

2100

Mohan Reddy (I) and Carpenter (CE). **Determination of metmyoglobin reductase activity in bovine skeletal muscles.** *Journal of Food Science* 56(5): 1991: 1161-1164

2101

Nuckles (RO), Smith (DM) and Merkel (RA). **Properties of heat-induced gels from beef skeletal, heart, lung and spleen protein fractions.** *Journal of Food Science* 56(5): 1991: 1165-1170

2102

Wild (JL), Sebranek (JG) and Olson (DG). **Grinding time and pressure developed in beef and pork: Effects of temperature and fat.** *Journal of Food Science* 56(5): 1991: 1171-1175

2103

Horgan (DJ), Kurth (LB) and Kuypers (R). **pH effect on thermal transition temperature of collagen.** *Journal of Food Science* 56(5): 1991: 1203-1204, 1208

Differential scanning calorimetry (DSC) was used to measure the thermal transition temp. of bovine intramuscular and tendon collagen. Equilibration of the collagen in buffers in the pH range 4.25-7.40 resulted in a decline in transition temp. with decreasing pH. This was more pronounced in collagen with higher concn. of aldimine crosslinks. It was proposed that the postmortem pH decline was responsible for the thermal transition temp. decline observed by other workers. Therefore, thermal transition temp. measurements must be performed on collagen samples that have been thoroughly



equilibrated to a common pH if they are to be used as indices of structural changes. AS

## **Pork**

2104

Brewer (MS), McKeith (F), Martin (SE), Dallmier (AW) and Meyer (J). **Sodium lactate effects on shelf-life, sensory and physical characteristics of fresh pork sausage.** *Journal of Food Science* 56(5): 1991: 1176-1178

2105

Decker (EA) and Crum (AD). **Inhibition of oxidative rancidity in salted ground pork by carnosine.** *Journal of Food Science* 56(5): 1991: 1179-1181

2106

Andersen (HJ) and Skibsted (LH). **Oxidative stability of frozen pork patties. Effects of light and added salt.** *Journal of Food Science* 56(5): 1991: 1182-1184

2107

Defreitas (Z) and Molins (RA). **Mechanically deboned pork use in fermented meat spreads.** *Journal of Food Science* 56(5): 1991: 1185-1190

## **Poultry**

### **Chickens**

2108

Zubillaga (MP) and Maerker (G). **Quantification of three cholesterol oxidation products in raw meat and chicken.** *Journal of Food Science* 56(5): 1991: 1194-1196, 1202

The results indicates that cholesterol oxidation products in meat and chicken could be measured at levels below 1 p.p.m. and 7-ketocholesterol and the isomeric 5,6-epoxides occur in such tissues with the amounts of the ketone exceeding those of the epoxides. SRA

## **Products**

### **Eggs**

2109

Satyanarayana Rao (TS). **Changes in solubility,  $\beta$ -carotene and development of non-enzymatic browning of spray-dried, foam-mat-dried and freeze-dried whole egg powders packed in different packaging materials.** *Journal of Food Science and Technology (India)* 29(4): 1992: 231-234

Drying conditions and packaging materials did not significantly influence solubility, destruction of  $\beta$ -carotene and development of non-enzymatic browning of spray-dried, foam-mat-dried and freeze-dried egg powders during storage at 4, 10 - 27 and 37 C upto 365 days. However, the storage at 42 and 55 C had significant effect on these attributes in all the 3 types of egg powders packed in cans and flexible pouches. AS

2110

Juhi Raikhy and Bawa (AS). **Studies on pickled chicken eggs.** *Journal of Food Science and Technology (India)* 29(4): 1992: 246-248

Sensory evaluation of egg pickles prepared with citric, acetic, lactic and tartaric acids indicated overall acceptability of the product made with lactic and tartaric acids when used alongwith 10% common salt, spices, condiments and oil. The storage period did not affect the sensory parameters. AS

2111

Wang (C) and Shelef (LA). **Factors contributing to antilisterial effects of raw egg albumen.** *Journal of Food Science* 56(5): 1991: 1251-1254

Sensitivity of *Listeria monocytogenes* strain Scott A and Brie-1 to several factors in raw egg albumen was investigated. A concn. of ca 15% of albumen in trypticase soy broth was listeristatic after 24 h at 35 C, and listericidal effects were observed at higher concn. Supplementation of albumen with Fe and biotin did not reverse the inhibition. Preheating of albumen (50 - 80 C) caused progressive loss of antilisterial effects. Supplementation of broth with lysozyme (> 1 mg/mL) produced antilisterial effects that were enhanced at pH 9; conalbumin (> 6 mg/mL) suppressed cell growth, while ovomucoid (> 2 mg/mL) was inhibitory only at pH 9. Results inferred that antilisterial effects of albumen were caused primarily by lysozyme and were enhanced by ovomucoid, conalbumin and alkaline pH. AS

2112

Chung (SL) and Ferrier (LK). **Partial lipid extraction of egg yolk powder: Effects on emulsifying properties and soluble protein fraction.** *Journal of Food Science* 56(5): 1991: 1255-1258

Solvent extraction to remove lipid from egg yolk decreased emulsifying properties of the resulting protein concentrates. Hexane and isopropanol had the least effects. Hexane-isopropanol extraction (77:23, w:w) reduced emulsifying activity (per 25 mg



protein) about 14% and protein solubility decreased from 15.7 to 13.8% but removed more than 50% of the lipid from egg yolk powder. Chloroform-methanol decreased total emulsifying activity upto 41%. Ovalbumin in the water-soluble fraction of extracted egg yolk powder reduced its emulsifying effectiveness. SRA

2113

Chung (SL) and Ferrier (LK). **Conditions affecting emulsifying properties of egg yolk phosvitin.** *Journal of Food Science* 56(5); 1991: 1259-1262

The effects of protein concn. (0.1 - 2.0%), oil vol. fraction (0.17 - 0.67), mixing speed (10,000 - 22,000 r.p.m.) and mixing time (0.5 - 8 min) on the emulsifying properties of phosvitin and bovine serum albumin (BSA) were compared. Emulsifying activity and emulsion stability increased with protein concn., oil vol. fraction and mixing. Effects of these variables were assessed quantitatively using an empirical equation. Mixing speed had the greatest influence and protein concn. had the least influence on emulsifying activity for both phosvitin and BSA. For emulsion stability, mixing speed had the greatest influence for phosvitin; oil vol. fraction had the greatest influence for BSA. Phosvitin was a better emulsifier than BSA at pH 7. AS

2114

Singh (RP) and Panda (B). **Storage stability of oil-based pickled quail eggs.** *Indian Journal of Poultry Science* 26(4); 1991: 221-225

Storage stability of quail eggs pickled in oil-based gravy containing 3% common salt and 2% acetic acid and stored in rigid (glass and plastic) jars or in flexible (PFP laminate, HDPE and PP) pouches for 9 and 15 months at mean ambient (28 C; 61% RH) and refrigerated (4 - 6 C; 80 - 85% RH) temp., were investigated. The rates of diffusion of both acid and salt were faster in egg white than in yolk and under ambient than refrigerated storage. Pickling caused approx. 2-fold increase in shear values of egg white. A progressive increase in % free fatty acids and a decrease in sensory quality occurred with storage time. Halophilic counts remained fairly low. Coliforms, anaerobes, *Salmonella* and coagulase-positive staphylococci were not detected in pickles throughout storage. The product could be safely stored in HDPE (330 G) pouches for 8 and 15 months under ambient and refrigerated conditions, respectively, instead of conventional rigid containers or intermediate cost PFP laminate. AS

2115

Maiti (D) and Biswas (S). **Effect of feeding deoiled niger (*Guizotia abyssinica*) cake on egg**

**production and egg quality.** *Indian Journal of Poultry Science* 26(4); 1991: 229-232

Deoiled niger cake (DNC) was included in the test diet of egg replacing groundnut cake (DGNC) protein at 0, 25, 50, 75 and 100% levels on equal nitrogenous and isocaloric basis. The birds were fed from 127 days of age. For egg quality studies, fresh eggs (8 eggs from each group) were collected at random at 278, 279 and 280 days of age. Egg wt., shape index, thin white %, albumin index, yolk index, shell thickness and Haugh unit score were recorded. When DNC was substituted up to 75% protein level, no adverse effect on egg production, egg wt., feed consumption per dozen of eggs and feed consumption per kg egg wt. was observed. Egg quality was also not affected by the substitution up to 100% DNC level except lowering of the egg wt. GS

2116

Singh (SV) and Panda (PC). **Peeling property of boiled eggs. A review.** *Poultry Guide* 29(6); 1992: 69-71, 73-79

Peeling property of boiled eggs is related to coagulation of egg white. The factors influencing coagulation of egg white are temp., dilution, salt, sugar, acid and alkali. The role of these are described. Storage, pH, shell treatment, oil coating, and the sp. of origin of egg are also factors influencing the peeling property of eggs. Cooking techniques which influence the peeling property are illustrated. The review concluded with the quality of cooked eggs including flavour, appearance and acceptability of peeled eggs. GS

## SEAFOODS

2117

Paturkar (AM), Sherikar (AA), Jayarao (BM) and. **Prevalence of *Salmonella* in meats and seafoods of Bombay city.** *Journal of Food Science and Technology (India)* 29(4); 1992: 242-243

Fifteen of the 166(9%) meat samples tested were positive for *Salmonella* in contrast to its absence in 96 fish samples, collected in Bombay. The serotype *S. saintpaul* predominated in the slaughterhouse meat as against the dominance of *S. mbandaka*, *S. adelaide*, *S. liverpool*, *S. anatum*, *S. derby* and *S. butantan* in market meat samples. *S. mbandaka* was frequently isolated from the retail market beef samples. AS

2118

Chung (KH) and Lee (CM). **Water binding and ingredient dispersion pattern effects on surimi**



**gel texture.** *Journal of Food Science* 56(5): 1991: 1263-1266

Water binding, thermotransitional properties and the dispersion pattern of ingredients were studied in relation to their texture-modifying effects in surimi gels. The DSC-measured bound water (unfreezable at -30 C) of surimi gels prepared with potato starch with and without pregelatinization highly correlated with compressive force ( $r = 0.94$ ) and inversely correlated with expressible moisture ( $R = 0.99$ ) of the gels. For nonfish proteins, however, physically bound water contributed more than the DSC-measured bound water to gel strength. The ability to undergo thermal transition (size expansion and water absorption) and dispersion pattern of ingredients were responsible for the differences in texture-modifying effects. AS

### Oysters

2119

Chai (T), Liang (KT), Pace (J) and Schlimme (DV). **Effect of heat processing on quality of pasteurized oysters.** *Journal of Food Science* 56(5): 1991: 1292-1294

Constant heating time at different temp. and constant temp. with different heating times combinations were studied to determine optimum heat process for oyster pasteurization in plastic pouches. Heating for 8 min at 75 - 76 C gave optimum physical and sensory quality to the product. Crude amylase and peroxidase extracts from oysters were inactivated by optimum treatment, but lipase retained about 15% of activity. AS

### Seal

2120

Synowiecki (J) and Shahidi (F). **Lipid and pigment extraction from mechanically separated seal meat.** *Journal of Food Science* 56(5): 1991: 1295-1297

Mechanically separated seal meat (MSSM) was washed 3x with water. Total hemoprotein pigment content of MSSM was reduced by 49.5% after the first washing, 65.1% after the second, and 67.5% after the third. Hunter L value (lightness) was increased from 14.8 to 32.4 after three washings. Hunter L values correlated ( $r = -0.990$ ) well with total hemoprotein content of the meat. The total lipid content of the meat was also reduced by 30.7 to 57.5%, depending on the original fat content of the sample after washings. The yield of meat protein after the second washing was increased from 32%

to 70.5% when a centrifugation instead of filtration method was employed. AS

### Squids

2121

Selvaraj (P), Indra Jasmine (G) and Jeyachandran (P). **Effect of polyphosphate dip treatment on frozen storage of Indian squid *Loligo duvauceli* Orbnig.** *Journal of Food Science and Technology (India)* 29(4): 1992: 248-249

Squid fillets (Mantles) given a dip treatment in aqueous trisodium polyphosphate sol. (5% w/v) prior to freezing showed reduced protein denaturation as evidenced by improved texture, decreased thaw and wt. loss, retention of salt soluble and  $\alpha$ -amino nitrogens as compared with control squids during frozen storage. The overall acceptance is better in polyphosphate treated samples than in control. AS

### Fish

2122

Voldrich (M), Dobias (J), Kalac (PJr) and Curda (D). **Changes of fatty acid composition during the processing of fish.** *Die Nahrung* 35(6): 1991: 663-664

Marked changes of the fatty acid comp. were observed in the smoked and marinated (processed) samples of mackerel meat. The processed fillets contained about 70% of n-3 polyunsaturated fatty acids (PUFAs). In case of smoked fish n-3 PUFAs loss was higher probably due to hydrolysis of fat before processing. In spite of the losses observed in the fish products-both marinated and smoked are an important source of n-3 PUFAs in human nutrition. BV

2123

Boyle (JL), Lindsay (RC) and Stuibler (DA). **Adenine nucleotide degradation in modified atmosphere chill-stored fresh fish.** *Journal of Food Science* 56(5): 1991: 1267-1270

Effects of CO<sub>2</sub> modified atm. on degradation of adenine nucleotides in chill-stored whitefish (*Coregonus clupeaformis*) and rainbow trout (*Salmo gairdneri*) were studied. K values were determined during storage up to 26 days. Results indicated CO<sub>2</sub> atm. did not alter K values compared to those observed for aerobically held fish. However, both CO<sub>2</sub> atm. and potassium sorbate-dipping of fillets caused decreases in hypoxanthine concn. compared to untreated aerobically held samples. AS



2124

Ohashi (E), Takao (Y), Fujita (T), Shimizu (Y) and Egashira (M). **Semiconductive trimethylamine gas sensor for detecting fish freshness.** *Journal of Food Science* 56(5): 1991: 1275-1279, 1286

Characteristics were studied on  $\text{In}_2\text{O}_3$  treated with 5 mol% MgO responses to 300 p.p.m. trimethylamine (TMA), dimethylamine (DMA), ammonia and other gases evolved from several different fish muscles stored under different conditions. Sensitivities of the  $\text{In}_2\text{O}_3$ -MgO sensor element were higher in order of TMA, DMA and ammonia over the operating range 330-620 C. Sensor responses were directly proportional to concn. of TMA in fish muscle in the presence of other evolved gases, irrespective of types of fish muscle or storage time. AS

### Finfish

2125

Himelbloom (BH), Brown (EK) and Lee (JS). **Microorganisms on commercially processed Alaskan finfish.** *Journal of Food Science* 56(5): 1991: 1279-1281

Whole and processed Alaskan fish were examined for aerobic plate counts, total coliforms, and *Escherichia coli*. Washing whole salmon and halibut reduced skin microbial counts from  $10^3/\text{cm}^2$  to  $10^2/\text{cm}^2$ . Whole and dressed fish had coliform counts  $< 13/\text{cm}^2$  and *E. coli* counts  $< 0.3/\text{cm}^2$ . Alaska pollock and Pacific cod fillets had microbial counts between  $10^3/\text{g}$  and  $10^6/\text{g}$ , coliform counts  $< 70/\text{g}$ , and *E. coli* counts  $< 4/\text{g}$ . Whole fish contained microbial flora predominated by *Moraxella* sp. whereas dressed fish and fillets had microbial flora consisting of *Arthrobacter/Corynebacterium*, *Flavobacterium*, and *Pseudomonas* sp. Microorganisms on conveyor belts and other contact surfaces may have contributed contaminants to fillets. AS

### Salmon

2126

Girard (B) and Nakai (S). **Static headspace gas chromatographic method for volatiles in canned salmon.** *Journal of Food Science* 56(5): 1991: 1271-1274

Amount of fish placed in vials as well as incubation temp. and time were factors studied to improve a static headspace sampling method applied to analyze volatiles in canned salmon. A random-centroid optimization (RCO) program was used to simultaneously search for the optimal levels

of other factors, namely, initial oven temp., column headpressure, and total flowrate. RCO was an effective optimization program while allowing testing of several treatments at a time. The optimal conditions of operation permitted detection of 80 volatile compounds, 34 of which were identified including aldehydes, alkanes, aromatic compounds, sulphur-containing compounds, alkenes, ketones, several miscellaneous compounds plus an alcohol and an acid. AS

2127

Durance (TD) and Collins (LS). **Quality enhancement of sexually mature chum salmon *Oncorhynchus keta* in retort pouches.** *Journal of Food Science* 56(5): 1991: 1282-1286

Sexually mature or late-run chum salmon muscle was packaged in thin-profile retort pouches and in conventional 307 x 200 metal cans. After retort processing to equivalent lethality, samples were evaluated by sensory and instrumental methods. The pouch product was firmer, more fibrous, drier and chewier than the canned. Less late-run flavour and greater overall acceptability were reported for pouch-processed samples. More free liquid was found in pouches than in cans. Histological examination revealed a more compact structure in chum muscle processed in retort pouches, possibly due to air over-pressure applied to flexible pouches during processing. AS

### Products

### Fish

2128

Konstance (RP). **Axial compression properties of kamaboko.** *Journal of Food Science* 56(5): 1991: 1287-1291

Kamaboko, a gelled seafood product from frozen surimi, has distinctive textural properties. Characterization of those properties, using an integrated approach to rheological studies, was accomplished by means of an instrumental texture profile analysis and evaluation of resultant stress-strain relationships. The material had near-ideal area expansion even at compressions of 60% while retaining its highly elastic texture. Apparently the product did not yield through 80% compression. Hardness of the kamaboko at 80% compression was characterized by a local max. at 37.5 C which may have been related to processing temp. of the initial surimi gel-set used in a double-gel-set procedure. Evaluation of stress-strain relationships confirmed the incompressible nature of the gel and showed relatively slight variations between the Young's



modulus and the deformability modulus. The elastic limit of the kamaboko increased significantly as temp. increased from 25 to 50 C. AS

## Fish oil

2129

Hsieh (Y-TL) and Regenstein (JM). **Factors affecting quality of fish oil mayonnaise.** *Journal of Food Science* 56(5); 1991; 1298-1301, 1307

Packaging conditions, antioxidants, metals and storage temp. affected the quality of fish oil (FO) mayonnaise (FOM). TBHQ was a successful antioxidant but ascorbate was not. N<sub>2</sub> retarded oxidation more than TBHQ. The metal content of FO was as low as soy oil (SO), suggesting that differences in quality deterioration were not due to metal content. FOM was more stable at refrigerator temp. (2 C) than at higher temp. (30 C). The sensory quality of deodorized FOM was equal to SO mayonnaise using TBHQ (0.02%), N<sub>2</sub>, and storing at 2 C for 14 wk. AS

## PROTEIN FOODS

2130

Ngo Som (J), Prajwala Mouliswar, Daniel (VA), Malleshi (NG) and Venkat Rao (S). **Digestibility of protein and starch in malted weaning foods.** *Journal of Food Science and Technology (India)* 29(4); 1992; 262-263

Roller-drying reduced *in vitro* protein digestibility of weaning foods based on malted/roasted maize/rice and malted cowpea. Cooking of weaning foods improved protein digestibility as compared to uncooked blends. *In vitro* starch digestibility was similar in various blends of cooked weaning foods. Increasing the malted material in weaning foods improved the digestibility of protein, but not the digestibility of starch. AS

## ALCOHOLIC AND NON-ALCOHOLIC BEVERAGES

### Non-alcoholic beverages

#### Coffee

2131

Kesavan (PC). **Protection by caffeine against oxic radiation damage and chemical carcinogens: Mechanistic considerations.** *Current Science* 62(12); 1992; 791-797

Caffeine administered after exposure to UV light enhances the damage to cells and organisms by inhibiting photoreactivation, excision and/or recombinational repair. But, if caffeine is already present in the system, it affords remarkable protection not only against O<sub>2</sub>-dependent component of radiation damage, but also against chemical carcinogens that require metabolic activation. The possible mechanistic aspects are discussed in this review. KAR

### Fruit juices

2132

Ganesh Kumar (C), Kanawjia (SK), Ladkani (BG) and Singh (S). **Recent advances in the processing of fruit juices.** *Indian Dairyman* 44(4); 1992; 167-179

Biotechnological techniques are increasingly used to get fruit juices which resemble the fresh juice in flavour, aroma, appearance and mouthfeel. Cost efficiency in processing methods; use of enzymes like pectinase, cellulase and hemicellulase in juice processing; nutritional evaluation of processed product; and development of new blended and nutrient fortified beverages with therapeutic value are the aspects considered. 61 references. GS

### Apple juices

#### Apple juice concentrates

2133

Dar (H), Zargar (MY) and Shah (GH). **Effect of processing operations and heat treatment on physico-chemical characteristics and microbiological load of apple juice concentrate.** *Indian Food Packer* 46(1); 1992; 45-50

Apple juice at various stages of processing did not show change in the pH, but acidity increased while concentrating to 71°Brix. Heat treatment of juice at 70 C for 1 h slightly increased total soluble solids due to the evaporation during this process. With the successive operations in concn. clarity increased after enzyme and gelation treatment. The raw juice was dominated by molds followed by yeasts whereas after processing, the yeasts dominated than molds and bacteria. Filtration decreased mold population. Heat treatment at 70 C for 1 h caused a net reduction to 0.68 - 4.6% in bacterial population, 1.03 - 3.23% in yeasts count and 1.19 - 2.44% in mold population, and an overall reduction of 96.9 - 98.8% in total microbial load. In the concn. the bacteria, yeasts and molds were within the range of 0.1/g. Osmophiles were < 100/g in unheated juice and were 1/g in conc. juice. GS



2134

Joshi (VK), Sandhu (DK), Attri (BL) and Walla (RK). **Cider preparation from apple juice concentrate and its consumer acceptability.** *Indian Journal of Horticulture* 48(4): 1991: 321-327

Fortification of apple juice concentrate with diammonium hydrogen phosphate as nitrogen source is essential for rapid fermentation. Must prepared by diluting the concentrate fermented faster than that diluted and ameliorated with sugar. The product prepared from direct dilution of the concentrate was found superior in sensory quality. SRA

### Grape juices

2135

Masoodi (FA), Kaur (B) and Kaur (H). **Effect of harvesting dates on the physico-chemical composition and quality of grape juice cv. Perlette.** *Indian Journal of Horticulture* 48(4): 1991: 328-333

Perlette var. grapes cultivated widely in Punjab matures only when the rainy season starts and the juice extracted is not of good quality. Hence, as assessment was made of the quality of juice extracted from Perlette var. grape harvested on 5 different dates starting from 1st to 18th June 1990 and also blending the juice with that extracted from Thompson-Seedless grapes in different proportions. Partial neutralization of the juice of early maturity, blending of Perlette-Thompson seedless grape juice up to 60:40 ratio, and addition of sugar up to 5% improved the early harvested Perlette grape juice quality. SRA

### Orange juices

2136

Nisperos-Carriedo (MO), Shaw (PE) and Baldwin (EA). **Changes in volatile flavour components of pineapple orange juice as influenced by the application of lipid and composite films.** *Journal of Agricultural and Food Chemistry* 38(6): 1990: 1382-1387

2137

Kimball (DA) and Norman (SI). **Processing effects during commercial debittering of California navel orange juice.** *Journal of Agricultural and Food Chemistry* 38(6): 1990: 1369-1400

2138

Baker (RA), Crandall (PG), Davis (KC) and Wicker (L). **Calcium supplementation and processing**

**variable effects on orange juice quality.** *Journal of Food Science* 56(5): 1991: 1369-1371

Untreated, pectinesterase (PE) treated, low pulp, and pectinase treated orange juices were fortified to 20% RDA of Ca with  $\text{CaCO}_3$  or  $\text{CaPO}_4/\text{lactate}$  (75/25). Neither Ca supplement adversely affected flavour, cloud density, settling pulp, or viscosity in untreated, low pulp, or pectinase treated juices. PE exposed orange juice held 4 h before pasteurization and Ca supplementation had less cloudy density, increased viscosity and more settling pulp.  $\text{CaCO}_3$  fortified PE exposed and pectinase treated juices had lower flavour scores, while  $\text{CaPO}_4$  suspensions caused slightly lower colour scores of all juices. AS

### Orange juice concentrates

2139

Crandall (PG) and Davis (KC). **Viscosity reduction and reformation of structure in orange concentrate as affected by homogenization within commercial taste evaporators.** *Journal of Food Science* 56(5): 1991: 1360-1364

Exp. were undertaken to determine if homogenization would reduce viscosity of bulk orange concentrate during commercial production and if a structure built-up during storage. Homogenizers, installed within two taste evaporators at separate citrus processing plants, were cycled "on" for homogenized samples and "off" for controls. In four replicates, homogenization significantly reduced viscosity to 19% average. Samples were stored for 6 months under tank farm conditions. Structure increased in 5 of 8 samples, but homogenized samples remained significantly lower in viscosity. This confirmed earlier pilot plant results under commercial operations. AS

### Passion fruit juices

2140

Shoseyov (O), Bravdo (BA), Siegel (D), Goldman (A), Cohen (S), Shoseyov (L), Ikan (R). **Immobilized endo- $\beta$ -glucosidase enriches flavour of wine and passion fruit juice.** *Journal of Agricultural and Food Chemistry* 38(6): 1990: 1387-1390

## FATS AND OILS

### Fats

2141

Sridhar (R) and Lakshminarayana (G). **Triacylglycerol compositions of some vegetable**



**fats with potential for preparation of cocoa butter equivalents by high performance liquid chromatography.** *Journal of the Oil Technologists Association of India* 23(3); 1991: 42-43

The triacylglycerols (TAG) composition of kokum (*Garcinia indica*), dhupa (*Vateria indica*), sal (*Shorea robusta*), mango (*Mangifera indica*), cocoa (*Theobroma cacao*) and papaya (*Carica papaya*) fats were determined by reversed phase HPLC. Kokum, dhupa, sal and mango TAG contained 85, 72, 76 and 62% of monounsaturated-disaturated glycerides. The papaya TAG contained tri-unsaturated and monounsaturated-diunsaturated glycerides as the major components. The TAG compositions of the indigenous vegetable fats could be useful in their modification into cocoa butter and in the identification of their presence when blended with cocoa butter. SRA

### **Vanaspati**

2142

**Shenolikar (I). Nickel content in vanaspati.** *Journal of the Oil Technologists Association of India* 23(3); 1991: 48-49

Ni content of vanaspati (hydrogenated fat) collected from market and household was analysed. The mean Ni content of the 25 samples was 1.55 p.p.m. with a range from 0.3 to 4.54 p.p.m. Vanaspati sold in sealed container or as loose did not show any difference in Ni content. Hence, vanaspati contributes negligible amount of Ni to the Indian diet. SRA

### **Oils**

2143

**Snyder (JM), Mounts (TL) and Holloway (RK). An analysis scheme for estimation of crude oil quality.** *Journal of the American Oil Chemist's Society* 68(5); 1991: 285-288

A seed analysis scheme was designed to rapidly estimate the quality of extracted oil. Factors of crude oil quality evaluated were: free fatty acids, oxidative status (Totox value), colour, and phosphatides (soybean) or wax (sunflower). Soybean and sunflower seeds subjected to extended storage at varying moisture contents were sampled at incremental time periods to yield fifty storage-damaged samples of each oilseed. Oil was extracted from 50 g lots of each samples and analysed for the crude oil quality factors according to standard methods. Alternative instrumental and chemical analyses of the quality factors were correlated with the standard methods. Hexanal

content, measured by headspace-gas chromatographic analysis of the ground full-fat meal, was correlated to the oxidative status. Crude oils recovered by rapid extraction, using sonication, and desolventation were monitored by spectrophotometry for colour correlation. Free fatty acid content was determined by titration methods and monitored by spectrophotometry. Modified turbidimetric methods estimated the phosphatides (soybean) or wax (sunflower seed) contents. The analysis scheme provides for the rapid estimation of oil quality as impacted by various pre- and post-harvest events that cause deterioration of oilseeds. AS

2144

**Sebedio (JL), Kaltaranta (J), Grandgirard (A) and Malkki (Y). Quality assessment of industrial prefried french fries.** *Journal of the American Oil Chemist's Society* 68(5); 1991: 299-302

An industrial production of prefried french fries using palm oil as a frying medium was studied over a period of 12 days. Samples of oil and french fries were withdrawn once a day. The quality of both the oil and the french fries was assessed using two types of tests. Some tests, such as the detn. of free fatty acid (FFA) and the detn. of thiobarbituric acid value (TBA), oxidfritest and Food Oil Sensor correspond to what was used by a quality control lab. More elaborate techniques such as the detn. of polar components, polymers and cyclic fatty acid monomers (CFAM) were also used. Only small increases of FFA, TBA, polar components and polymers were observed. However, in the case of palm oil, which contains a high percentage of diglycerides, it is more reliable to determine the quality of the oil using the amount of polymers instead of polar components which may include some diglycerides. Thus a high "polar components" value (up to 20-25%) would not necessarily reflect an altered sample. The max. amount of CFAM detected was 0.1% and they did not seem to be preferentially adsorbed on the french fries. These results, along with the sensory evaluations, showed that the french fries obtained in these production conditions were of good quality as far as the fat was concerned. AS

2145

**Adlof (RO), Rakoff (H) and Emken (EA). Preparation of deuterium-labelled methyl linoleate and its geometric isomers from natural seed oils.** *Journal of the American Oil Chemist's Society* 68(5); 1991: 303-306

Multi-gram quantities of deuterium-labelled methyl linoleate (methyl *cis*-9,*cis*-12-octadecadienoate) and its geometric isomers are readily synthesized from



**Crepis alpina** (70-80% 12, 13-epoxy-*cis*-9-octadeconic acid) seed oils. Methyl *cis*-9,*cis*-12-*trans*-9,*cis*-12-octadecadienoate-12,13- $d_2$  were prepared by the Lindlar-catalyzed reduction (with  $D_2$  gas) of methyl *cis*-9- and *trans*-9-octadecen-12-ynoates, respectively. Methyl *trans*-9-octadecen-12-ynoate was synthesized by the *p*-toluene-sulphinic acid-catalyzed isomerization of the corresponding *cis* isomer. Methyl *cis*-9,*cis*-12; *trans*-9,*cis*-12; *cis*-9,*trans*-12- and *trans*-9, *trans*-12-octadecadienoate- $d_2$ , - $d_4$ - and  $d_6$  were prepared by the Wittig coupling (with stereochemical control) of the appropriate  $d_2$ -,  $d_4$ - or  $d_6$ -alkyltriphenyl-phosphonium salt with methyl 12-oxo-*cis*-9- or *trans*-9-dodecenoate (prepared by the para-periodic acid cleavage of methyl 12,13-dihydroxy-*cis*-9- or *trans*-9-octadecenoate). The *cis* dihydroxy ester was synthesized from *Vernonia galamensis* seed oil by acetolysis, saponification and then esterification. The *cis* dihydroxy ester was isomerized by nitric acid/sodium nitrite to the *trans* form and purified by silver resin chromatography. Isotopic purities ranged from 88% (for the  $d_6$  isomers) to 99% (for the  $d_2$  isomers). AS

#### Cottonseed oils

2146

Kapseu (C), Kayem (GJ), Balesdent (D) and Schuffenecker (L). **The viscosity of cottonseed oil, fractionation solvents and their solutions.** *Journal of the American Oil Chemist's Society* 68(2); 1991; 128-130

Cold fractionation of cottonseed oil is made difficult by the high viscosity of the oil. This study was aimed at demonstrating the effect of solvents on the viscosity of mixtures between 0 and 25 C with a view to facilitating the fractionation of refined cottonseed oil. The solvents used were acetone, methylethylketone, methylisobutylketone, hexane and heptane. Measurements of viscosity were carried out by means of a capillary viscometer. The ratio of the viscosity of cottonseed oil to that of pure solvents is of the order of 300. The viscosities of solutions of various ratios of solvent to oil (1/3, 1/1, 3/1) are between those of cottonseed oil and the pure solvents. The effect of the solvent/oil ratio overrides that of solvent nature. The effect of solvent in reducing the viscosity of cottonseed oil is by descending order: acetone, hexane, methylethylketone, heptane, methylisobutylketone. AS

#### Rice bran oils

2147

Adhikari (J) and Adhikari (S). **Semi-quantitative detection of rape-mustard oil in rice bran oil.** *Journal of the Oil Technologists Association of India* 23(3); 1991; 50-52

The % of erucic, eicosenoic and linolenic acids can detect semi-quantitatively the proportion of Indian rape-mustard oil when present in rice bran oil. Fatty acid analysis of 22 Indian rape/mustard oil drawn from different seed var. and market samples were carried out. At 10% level of admixture or adulteration, the % would be 5.0 for erucic acid (22:1), 0.7 for eicosenoic acid (20:1) and 1.7 for linolenic acid (18:3). At 20% level, these % would be 10.0, 1.4 and 2.7 respectively. SRA

#### Soybean oils

2148

Palaniappan (S) and Proctor (A). **Evaluation of soy oil lutein isotherms obtained with selected adsorbents in hexane miscellas.** *Journal of the American Oil Chemist's Society* 68(2); 1991; 79-82

Bleaching of soy oil hexane/miscellas by adsorption of the lutein was examined. Quantitative differences in the adsorption behaviour between a bleaching clay, alkaline rice hull ash and acid rice hull ash were measured by comparing the constants, *K* and *n*, of Freundlich isotherms. All the isotherms were dose-dependent with *K* being inversely related and *n* directly related, to adsorbent mass. Overall, the bleaching clay performed better than the ashes. Alkaline ash had a larger *K* value than acid ash, but the *n* value of acid ash was greater. Equations of lutein adsorption were derived based on initial concn. on lutein and the amount adsorbed/g. Logarithmic plots produced constants,  $K^1$  and  $n^1$ , which were the intercept and slope, respectively. As adsorbent dose increased,  $K^1$  decreased and  $n^1$  increased for each adsorbent studied. These findings may have applications in the processing of dilute miscellas where low temp., low viscosity oil refining is desired. AS

2149

Schnepf (M), Spencer (G) and Carlat (J). **Chemical and sensory characteristics of stored menhaden oil/soybean oil blends.** *Journal of the American Oil Chemist's Society* 68(5); 1991; 281-284

The purpose of this study was to determine the feasibility of increasing the consumption of dietary  $\omega$ -3 fatty acids by incorporating menhaden oil into a French-type salad dressing. Menhaden/soybean oil blends of 10, 20 and 30% menhaden oil (w/w) were used to prepare an emulsified French salad dressing. The oil blends and salad dressings were stored at 22 C in the dark for 20 wk. The fatty acid



profile, peroxide value, and anisidine value were determined. The salad dressings also were evaluated by a sensory panel for flavour, aroma and aftertaste. The  $\omega$ -3 fatty acids were stable over time under these storage conditions. Peroxide values rose slowly and consistently over time reaching higher values when more menhaden oil was added. Peroxide values were also higher in the oil blends which were stored with air in the headspace and not flushed with argon. Anisidine values also were higher with each addition of menhaden oil but did not change over time except for the 100% menhaden oil which was stored in air. After 8 wks the sensory panel rated the salad dressing which contained menhaden oil as lower than the ones which did not contain menhaden oil. While a significant amount of  $\omega$ -3 fatty acids may be incorporated into foods by the addition of menhaden oil, the development over time of off-flavours must be controlled. AS

2150

Covey (JE) and Wan (PJ). **Hydrogenation of oxidized soybean oil.** *Journal of the American Oil Chemist's Society* 68(5): 1991: 337-338

Portions of refined and bleached soybean oil were stored at various temp. for various lengths of time, then hydrogenated to 70 iodine value (IV) to find the effect of peroxides on the rate of hydrogenation and on characteristics of hydrogenated product. Samples were treated up to 3 wk at up to 65 C and provided samples with peroxide values (PV) of up to 358. All samples were analyzed, hydrogenated and reanalyzed. Peroxides affected the fatty acid composition as determined by gas chromatography, the calculated iodine value based on fatty acid composition, and rate of hydrogenation. Peroxides also affected the selectivity of hydrogenation and slope of the solids curve in hydrogenated products. AS

#### Sunflower oils

2151

Semwal (AD) and Arya (SS). **Storage stability of refined sunflower oil in tins and HDPE bottles.** *Journal of Food Science and Technology (India)* 29(4): 1992: 250-252

Refined sunflower oil remains stable for 2 yr at room temp. when stored in high density polyethylene (HDPE) bottles and sealed tins without development of perceptible off-flavour or odours. Peroxide, TBA, total carbonyls and anisidine values increased during storage and the changes correlated linearly with the storage period. Increases in these values were more in leaking containers. AS

#### SPICES AND CONDIMENTS

2152

Carbonell (ES). **Extraction of flavours with supercritical carbon dioxide.** *Cereal Foods World* 36(11): 1991: 935-937

Flavour extraction with supercritical CO<sub>2</sub> results in excellent flavour characteristics with other products such as spices because of the low temp. and inert atm. involved and the potential selection of specific aroma fractions, extractions of spices and hops, ranges of solubility for CO<sub>2</sub>, method of extraction, plant operations, examples of extraction of spices and flavours (ginger, pepper and vanilla) and dealcoholization of beverages are the aspects covered. BV

2153

Amita Srivastava and Jain (PC). **Seed mycoflora of some spices.** *Journal of Food Science and Technology (India)* 29(4): 1992: 228-230

Survey of fungal contaminants of bishopweed, black pepper, coriander and cumin from Sagar district of Madhya Pradesh, India showed occurrence of *Aspergillus flavus*, *A. niger*, *A. ochraceus*, *Penicillium*, spp., *Rhizopus arrhizus*, *R. stolonifer* and *Syncephalastrum racemosum*. However, their frequency varied widely. Samples of test spices yielded slightly lower fungal species with the use of spice extract agar medium as compared to malt-salt agar. AS

#### Pepper

2154

Gopalakrishnan (N) and Thomas (PP). **Studies on colour retention in pepper subjected to different treatments.** *Journal of Food Science and Technology (India)* 29(4): 1992: 256-257

Fresh pepper (*Piper nigrum* L.) was subjected to treatments such as microwave exposure, microwave exposed boiling water blanching and direct boiling water blanching to study the extent of green colour retention. The variation in the chlorophyll content at different stages was insignificant among the samples subjected to the same treatment. The best colour retention was observed in the microwave exposed boiling water blanched samples. AS



2155

Kuchiba-Manabe (M), Matoba (T) and Hasegawa (K). **Sensory changes in umami taste of inosine 5'-monophosphate solution after heating.** *Journal of Food Science* 56(5): 1991: 1429-1432

Discrimination in umami taste of inosine 5'-monophosphate (IMP) sol. caused by thermal degradation was investigated by sensory evaluation. The difference threshold of umami taste of 0.005% IMP sol. in the presence of 0.05% monosodium glutamate was 0.002%. The difference threshold of a 0.005% IMP sol. decreased by about one half when heated at 95 C for 15 h. Inosine, one of the main products of the thermal degradation of IMP, had a bitter taste. The detection threshold of inosine varied widely among panelists. Heating a 0.005% IMP solution at 95 C for 15 h formed inosine at about one tenth of its lowest detection threshold. AS

2156

Nunes (RV), Rhim (JW) and Swartzel (KR). **Kinetic parameter evaluation with linearly increasing temperature profiles: Integral methods.** *Journal of Food Science* 56(5): 1991: 1433-1437

Two integral methods for kinetic parameter estimation with linear temp. profiles were tested using simulated and experimental data sets. They were the second rational approximation method (SRAM) and the equivalent point method (EPM) using least squares nonlinear regression (LSNR) and weighed least squares nonlinear regression (WLSNR). For three simulated data sets, the SRAM with WLSNR yielded accurate parameter estimation. For the experimental data set, both SRAM and EPM using WLSNR yielded accurate parameter estimation. The standard error in activation energy were 37.3% (for SRAM) and 46.7% (for EPM) lower than that of the differential method. The SRAM and WLSNR was the best parameter estimation procedure. AS

## FOOD STORAGE

Nil

## INFESTATION CONTROL AND PESTICIDES

Nil

Nil

## TOXICOLOGY

2157

Lewerenz (H-J), Mieth (G), Bleyl (DWR), Plass (R) and Elsner (A). **Toxicological evaluation of sucrose carbonic acid esters in subchronic feeding studies in rats. Part 2. Effect of sucroacetoglycerides.** *Die Nahrung* 35(6): 1991: 655-662 (De)

Groups of male and female rats received a sucroacetoglycerides containing product (SAG) for 3 months at dietary levels of 0, 2.5, 5 and 10%. Food consumption was initially increased in females at all SAG-levels. After two wks significant increases in food intake were observed in males and females fed 10% SAG throughout the feeding period. The serum analysis revealed significantly elevated activity in serum alkaline phosphatase at the highest SAG-level in males and females after 6 wks and in females after 13 wks. Histological changes related to the SAG-feeding were noted in the intestinal lymph nodes of male and female animals fed 10% SAG. The no-adverse effect level established in this subchronic feeding study was 5% SAG in the diet of rats, equivalent to a daily intake of 3.6 g/kg body wt. in males and 4.0 g/kg in females. AS

2158

Choudhary (DN), Sahay (GR) and Singh (JN). **Effect of some mycotoxins on reproduction in pregnant albino rats.** *Journal of Food Science and Technology (India)* 29(4): 1992: 264-265

Effect of oral administration of aflatoxin B<sub>1</sub>, patulin and kojic acid on reproductive performance of pregnant rats showed significant anti-implantational activity. Patulin and kojic acid exhibited significant abortifacient activity. A significant loss of viability among the litters was also noticed in mycotoxin fed rats except for those fed with patulin. AS

## FOOD LAWS AND REGULATIONS

Nil



# AUTHOR INDEX

- Aaron (DK)  
2095  
Abbott (J)  
2060  
Abo-ElNile (E)  
2010  
Acuff (GR)  
2096  
Addis (PB)  
2098  
Adhikari (J)  
2147  
Adhikari (S)  
2147  
Adlof (RO)  
2145  
Agarwal (AK)  
2048  
Agarwal (JKP)  
2058  
Agarwal (S)  
2058  
Agullera (JM)  
2064  
Aharoni (N)  
2028  
Ahire (CN)  
2045  
Akerman (M)  
2028  
Alem (F)  
2040  
Aly (ME)  
2073  
Amita Srivastava  
2153  
Amundson (CH)  
2072  
Andersen (HJ)  
2106  
Andrich (G)  
2027  
Angelini (P)  
2097  
Annapure (SG)  
1986  
Appu Rao (AG)  
2086  
Armagost (MS)  
1968  
Arora (DK)  
1968  
Arya (SS)  
1984 2151  
Attri (BL)  
2134  
Ayyad (K)  
2010  
Azuara (E)  
1959  
Badrie (N)  
2013 2014  
Bajaj (VD)  
2048  
Baker (RA)  
1967 2138  
Balasubrahmanyam (N)  
1958  
Baldwin (EA)  
2136  
Balesdent (D)  
2146  
Banerji (R)  
2002  
Banwari Lal  
2042  
Baranth (AM)  
2045  
Bartholomai (GB)  
2008  
Bawa (AS)  
2110  
Baxter (L)  
2016  
Bell (DA)  
2062  
Benson (ME)  
2001  
Beristain (CI)  
1959  
Bernardi Don (LS)  
2008  
Bhagya (S)  
1999  
Bhirud (VS)  
1974 1975  
Bhojaraj (SK)  
2044  
Bhupinder (K)  
2049  
Binder (RG)  
2001  
Bingham (C)  
2098  
Biswas (S)  
2115  
Bleyl (DWR)  
2157  
Bodhankar (SS)  
1986  
Boyle (JL)  
2123  
Bravdo (BA)  
2140  
Bressani (R)  
2019  
Brewer (MS)  
2104  
Brown (EK)  
2125  
Brown (WJ)  
2022  
Carbonell (ES)  
2023 2152  
Carlat (J)  
2149  
Carpenter (CE)  
2100  
Cha (G-S)  
1978  
Chadha (KL)  
2033  
Chai (T)  
2119  
Chambers (P)  
2084  
Chandan (RC)  
2075  
Chandrasekar (N)  
2052  
Changade (SP)  
2079  
Charjan (SKU)  
2004  
Chauhan (SS)  
2056  
Chavan (KD)  
2066  
Chimi (H)  
1979  
Chin (KB)  
2087  
Chlou (RY-Y)  
1997  
Chou (TC)  
2029  
Choudhary (DN)  
2158  
Chung (KH)  
2118  
Chung (SL)  
2112 2113  
Cillard (J)  
1979  
Cillard (P)  
1979  
Clark (PK)  
2005  
Cohen (S)  
2140



Colbert (LB)  
2080  
Collins (LS)  
2127  
Conway (WS)  
2026  
Costell (E)  
2023  
Covey (JE)  
2150  
Crandall (PG)  
2138 2139  
Cross (HR)  
2096  
Crouse (JD)  
2094  
Crum (AD)  
2105  
Curda (D)  
2122  
Daeschel (MA)  
2018  
Dallmiller (AW)  
2104  
Daniel (VA)  
2130  
Dar (H)  
2133  
Davidson (SD)  
1970  
Davis (KC).  
2138 2139  
Decker (EA)  
2080 2105  
Defreltas (Z)  
2107  
Desai (HK)  
2070  
Desai (NB)  
2070  
Dixit (AN)  
2046  
Doblas (J)  
2122  
Dodds (GT)  
2022  
Donnelly (WJ)  
2063  
Dubey (HK)  
2051  
Dubey (Y)  
2051  
Duran (L)  
2023  
Durance (TD)  
2127  
Economou (KD)  
1977  
Egashira (M)  
2124

El-Abbassy (MZ)  
2073  
El-Mahdy (AR)  
1995  
El-Makhzangy (A)  
2010  
Eldon Brown (G)  
2031  
Elsner (A)  
2157  
Emken (EA)  
2145  
Epley (RJ)  
2098  
Etlevant (P)  
2038  
Farkas (DF)  
1971  
Ferguson (DM)  
2093  
Ferrier (LK)  
2112 2113  
Florentini (R)  
2027  
Flath (RA)  
2001  
Fleming (HP)  
2018  
Frias (J)  
1992  
Fu (B)  
2069  
Fuchs (Y)  
2028  
Fujita (T)  
2124  
Galewska (E)  
1973  
Ganesh Kumar (C)  
2132  
Garcia (HS)  
1959 2072  
Garcia-Vela (LA)  
2019  
Gautam (GK)  
2048  
Gehlawat (JK)  
1957  
Giles (JE)  
2093.  
Girard (B)  
2126  
Goldman (A)  
2140  
Gopala Krishna (AG)  
1964  
Gopalakrishnan (N)  
2154  
Graham (HD)  
1966

Grandgirard (A)  
2144  
Griffin (VK)  
1985  
Gunasekaran (N)  
2052  
Gupta (RK)  
2002  
Gurugovind (J)  
2052  
Ha (JK)  
2074 2088  
Hamaker (BR)  
1985  
Hansen (AP)  
1968  
Haridas Rao (P)  
2061  
Hasegawa (K)  
2155  
Hayakawa (K-I)  
1960 1961  
Hernandez (E)  
1967  
Hill (CGJr)  
2072  
Himelbloom (BH)  
2125  
Hirano (J)  
1978  
Holloway (RK)  
2143  
Hoover (DG)  
1971  
Horgan (DJ)  
2103  
Hsieh (Y-TL)  
2129  
Huang (CJ)  
1989  
Hung (SC)  
2082  
Hung (Y-C)  
1998  
Idouraine (A)  
1991  
Ikan (R)  
2140  
Indoni (SB)  
1986  
Indra Jasmine (G)  
2121  
Indrani (D)  
1987  
Ingle (VK)  
1976  
Isoda (Y)  
1978  
Jabunathan (R)  
1990



Jain (PC)	Kilara (A)	Laxmi Narayan
2153	2081	2050
Jain (PF)	Kim (N-K)	Lee (BH)
2045	1998	2076
James (J)	Kim Ha (J)	Lee (CM)
1983	2092	2118
Jayarao (BM)	Kimball (DA)	Lee (JE)
2117	2137	1962
Jeyachandran (P)	Kinsella (JE)	Lee (JS)
2121	1965 2064	2125
Joshi (JT)	Kokini (JL)	Lesschaeve (I)
2045	2029	2038
Joshi (NS)	Konstance (RP)	Levin (RE)
2039	2128	2097
Joshi (VK)	Koohmarale (M)	Lewerenz (H-J)
2032 2134	2094	2157
Juhl Raikhy	Kouassi (MA)	Liang (KT)
2110	2087	2119
Kaltaranta (J)	Kramer (GF)	Libudzisz (Z)
2144	2026	1973
Kakuda (N)	Krol (B)	Lindsay (RC)
2015	2091	2074 2088 2092 2123
Kakuda (Y)	Kroll (J)	Lucia (LM)
2015	2000	2096
Kalac (PJr)	Kuchiba-Manabe (M)	Ludford (PM)
2122	2155	2022
Kallyamurthy (S)	Kulkarni (DN)	Madsen (M)
2054	2030	2084
Kaloti (SV)	Kulkarni (KD)	Maerker (G)
2079	2030	2108
Kanawjia (SK)	Kulkarni (MB)	Maiti (D)
2132	2066	2115
Kanda (K)	Kulkarni (SD)	Malkki (Y)
2059	2006	2144
Kang (JII)	Kumar (KR)	Malleshi (NG)
2009	1958	2130
Kanno (C)	Kumar (VU)	Malshe (VC)
2067	2065	2047
Kanwar (RS)	Kurth (LB)	Mangino (ME)
2043	2103	2081
Kapseu (C)	Kuypers (R)	Marlanchuk (M)
2146	2103	1996
Kapur (PPatitVL)	Labuza (TP)	Martin (SE)
2048	2069	2104
Kashima (M)	Ladkani (BG)	Masoodi (FA)
1978	2132	2135
Kaur (B)	Lakshminarayana (G)	Matoba (T)
2135	2141	2155
Kaur (H)	Langlois (BE)	Matsumura (Y)
2135	2095	2009
Kayem (GJ)	Langlois (D)	Mattila (M)
2146	2038	2037
Kesavan (PC)	Larrigaudiere (C)	McCollum (TG)
2131	2021	2031
Khanna (KR)	Latche (A)	McDonald (LC)
2002	2021	2018
Khanna (SK)	Lau (OL)	McDonald (RE)
1981	2025	2031 2034
Khorgade (AB)	Lauro (GJ)	McGuire (CF)
2079	1980	2060



McWatters (KH)

1993 1994

McKelth (F)

2104

Means (WJ)

2095

Mehta (NJ)

2041

Meir (S)

2028

Mellowes (WA)

2013 2014

Menichillo (DA)

2099

Merkel (RA)

2101

Meyer (J)

2104

Mieth (G)

2157

Miller (AJ)

2099

Miller (MF)

2094

Miller (RK)

2096

Miller (WR)

2031 2034

Milne (JAC)

2084

Minerich (PL)

2098

Mishra (HN)

2007

Mitchell (FG)

2035

Mitchell (GE)

2093

Miyazawa (T)

1978

Mohan Reddy (I)

2100

Moldenhauer (KAK)

1985

Mollins (RA)

2107

Moody (WG)

2095

Mori (T)

2009

Mounts (TL)

2143

Mukherjee (RK)

2007

Mulvihill (DM)

1985

Murthy (TSN)

2053

Nagarajan (L)

2065

Naldoo (RJ)

2093

Nakai (S)

2126

Nanos (GD)

2035

Nayital (RK)

2024

Ngo Som (J)

2130

Ngoddy (PO)

1993 1994

Nisperos-Carriedo (MO)

2136

Norman (SI)

2137

Nuckles (RO)

2101

Nunes (RV)

2156

O'Palka (J)

2060

Ohashi (E)

2124

Okechukwu (PE)

1993 1994

Olson (DG)

2102

Onkar Singh

2043

Oreopoulou (V)

1977

Owusu-Ansah (YJ)

1996

Paakkonen (K)

2037

Pai (J)

2119

Pal (SK)

2040

Palaniappan (S)

2148

Palombo (R)

2091

Panda (B)

2114

Panda (PC)

2116

Pandian (S)

2071

Papadopoulos (LS)

2096

Park (YW)

2087

Parmar (C)

2024

Patel (HS)

2041

Patel (MP)

2041

Patil (VK)

2030

Paturkar (AM)

2117

Pawar (SG)

2030

Payne (CA)

2095

Pech (JC)

2021

Pegg (RB)

2089 2090

Pilug (IJ)

1972

Phatak (RN)

2047

Philosoph-Hadas (S)

2028

Pilosof (AMR)

2008

Pintauro (N)

2029

Plass (R)

2157

Pomnier (SA)

2085

Poonam Aggarwal

2011

Prabhakara Raju (RV)

1969

Prajwala Mouliswar

2130

Prakash (GS)

2033

Prakash (V)

1999 2003

Prins (A)

2091

Proctor (A)

2148

Raghava Naidu (R)

1969

Rahmani (M)

1979

Rahmen (MH)

2040

Rakoff (H)

2145

Ramasamy (P)

2054

Ranganna (S)

2017

Rao (KSN)

2050

Rao (MA)

1993 1994

Razla Tasneem

2003

Reagan (JO)

2094



Rector (D)	Shah (US)	Singh (S)
1985	1983	2077 2132
Reddy (BMC)	Shahidi (F)	Singh (SV)
2033	2089 2090 2120	2116
Regenstein (JM)	Shamsuzzaman (K)	Sinha (R)
2129	2090	1988
Rhum (JW)	Sharada (D)	Skibsted (LH)
2156	1988	2106
Robinson (RK)	Sharma (GC)	Smith (DM)
2068	2051	2101
Rogers (SA)	Sharma (KD)	Snyder (HE)
2093	2032	2005
Sahay (GR)	Sharma (KP)	Snyder (JM)
2158	2049	2143
Sai Kumar (KS)	Sharma (KR)	Sommovigo (G)
2048	2036	2027
Saikia (L)	Sharma (PC)	Spencer (G)
2017	2032	2149
Saini (SPS)	Sharma (RS)	Sridhar (R)
2011	2039	2141
Sakal (N)	Sharp (JL)	Srinivas (K)
1961	2034	2050
Samona (A)	Shattuck (VI)	Srinivasan (KS)
2058	2015	1999
Sandberg (A-S)	Shaw (PE)	Srinivasan (S)
1982	2136	2052
Sandhu (DK)	Shelef (LA)	Steinke (LW)
2134	2111	2062
Sapp (RE)	Shelp (BJ)	Strange (ED)
1970	2015	2083
Satyanarayana Rao (TS)	Shenolikar (I)	Stulber (DA)
2109	2142	2123
Schlimme (DV)	Shere (DM)	Stutz (HK)
2119	2030	2097
Schnepf (M)	Sherikar (AA)	Styles (MF)
2149	2117	1971
Schooner (F)	Shimizu (Y)	Subrahmanyam (VVR)
2071	2124	1974 1975
Schuffenecker (L)	Shimomura (Y)	Subramanian (V)
2146	2067	1990
Sebedio (JL)	Shoseyov (L)	Sud (G)
2144	2140	2024
Sebranek (JG)	Shoseyov (O)	Sudhir Singh
2102	2140	1981
Seetharama (N)	Shukla (BK)	Suresh (P)
1990	2057	1976
Selvakumaran (R)	Shyu (S-L)	Svanberg (U)
2065	1997	1982
Selvaraj (P)	Siegel (D)	Swartzel (KR)
2121	2140	2156
Semwal (AD)	Silverman (GJ)	Syed (HM)
1984 2151	2097	1986
Sen (DC)	Simard (RE)	Synowiecki (J)
2078	2071 2076	2120
Shackelford (SD)	Singh (JN)	Taha (SN)
2094	2158	2073
Shah (CM)	Singh (RK)	Takano (E)
1963	1962	2067
Shah (GH)	Singh (RP)	Takao (Y)
2133	2114	2124



Tambat (RV)	2083	Waters (LJr)
2079	Van Roon (PS)	2016
Tan (LT)	2091	Weber (CW)
2093	Vanderzant (C)	1991
Taoukis (PS)	2096	Wei (TM)
2069	Vashi (RD)	2006
Tarar (JL)	2041	Wergin (WP)
2004	Vasudeva (TR)	2012
Tatsumi (Y)	2048	Wheeler (TL)
2012	Vatsala (CN)	2094
Tayagi (ID)	2061	Whipple (G)
2055	Venkat Rao (S)	2094
Teotla (MS)	2130	Wicker (L)
2020	Venkataraman (LV)	2138
Tewari (BD)	1956	Wijeratne (WB)
2077	Venkateswara Rao (G)	2006
Thakar (PN)	1987	Wild (JL)
2070	Venkateswara Rao (P)	2102
Thakur (NS)	1990	Wolfe (RR)
2036	Verma (VK)	1960
Thomas (PP)	2055	Yastremski (R)
2154	Vidal-Valverde (C)	2025
Thomopoulos (CD)	1992	Yasuda (Y)
1977	Vijayalakshmi (V)	2059
Thompson (MP)	1976	Yensen (SB)
2083	Viplin Kumar	1991
Tochikubo (K)	2077	Youssef (MM)
2059	Voldrich (M)	1995
Trepanier (G)	2122	Zargar (MY)
2076	Vyas (HK)	2133
Triantaphyllides (C)	1963	Zauberman (G)
2021	Walla (RK)	2028
Tsal (C-L)	2134	Zayas (JF)
1997	Wan (PJ)	1989 2082
Tsukada (T)	2150	Zechman (LG)
1961	Wang (C)	1972
Tuci (A)	2111	Zlena (HM)
2027	Wang (CY)	1995
Umesh Kumar (S)	2026	Zinnai (A)
2065	Wang (J)	2027
Valdya (SD)	1960	Zubillaga (MP)
1974 1975	Watada (AE)	2108
Van Hekken (D)	2012	



## SUBJECT INDEX

### Acceptability

- apple juice concentrate, acceptability & cider preparation from 2134
- beef top rounds, sodium lactate & acceptability of cooked roast stored 2096

### Acidification

- cheese manufacture & acidification 2075
- cucumber, acidification & microorganisms during fermentation of 2018

### Adenine nucleotide

- white fish/rainbow trout, adenine nucleotide degradation in modified atm. chill-stored 2123

### Ageing

- beef steaks, ageing & properties of 2093

### Albumin

- tepary bean albumin, functional properties of 1991

### Alkylphenol

- meat, alkylphenols & flavour of red 2088

### Amaranthus

- grain amaranth, protein fractions of 2019

### Amine

- Mucor miehei lipase & primary amines alkylation 1975

### Amino acid

- faba bean medamms, cooking temp./time & amino acid composition of cooked 1995

### Antibacterial property

- eugenol, antibacterial activity of 1976

### Antioxidant

- perilla oil, phospholipid antioxidant effect on 1978
- phenolic antioxidant, peroxy/hydroxyl radical scavenging activity of 1979
- plant extracts, antioxidant activity of 1977
- whey, antioxidant from acid 2080

### Apple

- polyamine & softening inhibition in Golden Delicious/McIntosh apple 2026
- respiration model for stored apples 2027
- shelf-life extension/bavistin/diphenylamine in apples 2024

- storage controlled/modified atm. & quality of Golden Delicious apples 2025

### Apple juice concentrate

- acceptability & cider preparation from apple juice concentrate 2134
- processing/heat treatment & physicochemical properties/microbial quality of apple juice concentrate 2133

### Arachin

- groundnuts, proteolytic enzymes & arachins of 1999

### Arthrobacter

- finfish, Arthrobacter in commercially processed Alaskan 2125

### Asparagine

- Bacillus subtilis spores, L-asparagine & germination response of 2059

### Asparagus

- spears, packaging & quality changes in stored 2016

### Aspergillus niger

- butter oil emulsions, Aspergillus niger lipase action on 2072

### Atomic absorption spectrometry

- food, Na microwave digestion AAS analysis in 1970

### Auramine

- toxicity of auramine 1981

### Avocado

- peels, ripening & lipid peroxidation products in avocado 2028

### Bacillus cereus

- milk, B. cereus & quality of low-temp. stored 2065

### Bacillus stearothermophilus

- spores, media & recovery of B. stearothermophilus 1972

### Bacillus subtilis

- spores, sugar/L-asparagine & germination response of B. subtilis 2059

### Bacteria

- beef, bacteria & spoilage of ground 2097
- soy protein concentrate, bacterial protease modification of 2008

### Basil

- antioxidant activity of basil 1977

### Bavistin

- apple, shelf-life extension bavistin in 2024

### Beef

- bacteria/volatile compounds & spoilage of ground beef 2097
- mixtures, properties of wild rice/ground beef 2098
- protein fractions, gels heat-induced properties of beef 2101
- sausages, blood & nitrite antibotulinal efficacy in beef 2099
- temp./fat & grinding time/pressure developed in beef 2102
- tenderness, prediction of beef 2094
- top rounds, sodium lactate & acceptability of cooked roast stored beef 2096

### Beef steak

- microbial quality of precooked restructured beef steaks 2095
- tenderization/ageing/thawing & properties of beef steaks 2093

### Beverage

- India, beverage production prospects in 1956
- pumpkin beverage 2020

### Bifidobacteria

- dairy products, bifidobacteria enumeration in 2068

### Biotechnology

- dairy processing, biotechnology applications in 2063

### Blanching

- pumpkins, blanching of 2020

### Blue VRS

- toxicity of blue VRS 1981

### Bottle

- sunflower oil, stability of refined HDPE bottle stored 2151

### Bovine

- milk fat emulsions, bovine milk fat globule membrane & properties of 2067
- skeletal muscles, metmyoglobin reductase activity detn. in bovine 2100

### Brassica

- chlorophyll loss during processing of Brassica 2011

### Bread

- fruit bread 2061
- soy flour fortified bread 2006

### Breadmaking

- wheat flour, milling method & breadmaking characteristics of



- 1987
- Butter**  
fatty acid flow  
injection/enzymatic analysis for  
butter 2071
- Butter oil**  
emulsions, *Aspergillus niger*  
lipase action on butter oil 2072
- Caffeine**  
oxic radiation damage/carcinogen  
control by caffeine, review 2131
- Cake**  
methylcellulose gum &  
structure/texture of  
microwave-baked cake 2062
- Calcium**  
orange juice, Ca & quality of 2138
- Calcium chloride**  
apple, shelf-life extension  
CaCl<sub>2</sub> in 2024
- Calorimetry**  
whey protein concentrates,  
thermal properties & solubility  
of 2081
- Candelilla**  
wax emulsion,  
preparation/stability of  
Candelilla 1967
- Canned food**  
salmon, volatile compounds  
GC detn. in canned 2126  
vegetables, thermal process for  
canned 2017
- Canning**  
Brassica, chlorophyll loss during  
canning of 2011  
pumpkins, canning of 2020
- Canola**  
microwave & myrosinase in canola  
seeds 1996
- Carbon dioxide**  
flavour, extraction supercritical  
CO<sub>2</sub> of 2152  
peaches, temp. & CO<sub>2</sub>  
injury/flesh softening in 2035
- Carcinogen**  
caffeine for carcinogen control,  
review 2131
- Carnosine**  
pork, carnosine & rancidity  
inhibition of salted 2105
- Carrot**  
stick, scanning electron  
microscopy of carrot 2012
- Carthamus tinctorius**  
see Safflower
- Casein**  
qualitative/quantitative detn. of  
caseins 2083
- Cassava**  
extrudates, extrusion variables  
effect on cassava 2013
- Cassava flour**  
extrudate, texture/microstructure  
of cassava flour 2014
- Cereal**  
India, cereal production  
prospects in 1956
- Cheddar cheese**  
*Lactobacillus casei* & accelerated  
maturation of Cheddar cheese 2076  
spray drying for accelerated  
ripened Cheddar cheese 2077
- Cheese**  
Italian cheese, fatty  
acids/phenolic compounds &  
flavour  
in aged 2074  
accelerated cheese ripening 2063  
lecithin & yield/quality of soft  
cheese 2073  
manufacture, acidification &  
cheese 2075
- Chemical property**  
beef steaks,  
tenderization/aging/thawing &  
chemical properties of 2093  
milk fat emulsions, bovine milk  
fat globule membrane & chemical  
properties of 2067  
wheat flour, milling method &  
chemical properties of 1987
- Chemistry**  
pumpkin, chemistry of, review 2020
- Chicken**  
cholesterol oxidation products in  
chickens 2108  
eggs-pickled 2110
- Chilling**  
grapefruit,  
thiabendazole/imazalil & chill  
injury/decay of 2031  
tomatoes, chilling & surface  
colour changes in 2022
- Chlorophyll**  
Brassica, chlorophyll loss during  
processing of 2011
- Chocolate**  
milk chocolate 2039
- Cholesterol**  
goat organ/muscle meat,  
cholesterol in 2087  
meat, cholesterol oxidation  
products in 2108
- Chymotrypsin**  
*Sesame indicum*, chymotrypsin &  
 $\alpha$ -globulin of 2003
- Cider**  
apple juice concentrate,  
acceptability & cider  
preparation from 2134
- Citrate**  
*Lactococcus lactis*, citrate  
metabolism in 1973
- Citrus**  
pectin, conformation of citrus  
2029
- Collagen**  
pH & thermal transition temp. of  
collagen 2103
- Colorimetry**  
butter, fatty acid colorimetric  
analysis for 2071
- Colour**  
meat,  $\gamma$ -irradiation &  
colour of nitrite-free cured 2090  
molasses, sugar colour & purity  
of 2054  
natural colour 1980  
pepper, microwave/blanching &  
colour retention in 2154  
pork, carnosine & rancidity  
inhibition of salted 2105  
tomatoes, chilling & surface  
colour changes in 2022  
toxicological evaluation of  
non-permitted food colours 1981
- Condensing**  
power reduction in condensing  
system 2045
- Container**  
pumpkin, containers for 2020
- Cooking**  
faba bean mediamis, cooking  
temp./time & amino acid  
composition/antinutritional  
factors of cooked 1995  
strawberry jam, cooking &  
volatile compounds in 2038
- Coriander**  
seed mycoflora of coriander 2153
- Corn**  
dried distiller's grains with  
solubles 2060  
roli quality & corn  
characteristics 1988
- Corn flour**  
emulsifying capacity/emulsion  
stability of corn germ protein  
flour 2082  
phenolic acid & taste of corn  
germ protein flour products 1989
- Corn product**  
phenolic acid & taste of corn  
germ protein flour products 1989
- Corn protein**  
emulsifying capacity/emulsion  
stability of corn germ protein  
flour 2082  
phenolic acid & taste of corn



- germ protein flour products 1989
- Corrosion**  
stainless steel, corrosion prevention of 1963
- Corynebacterium**  
finfish, corynebacterium in commercially processed Alaskan 2125
- Cottonseed oil**  
viscosity of cottonseed oil 2146
- Cowpea**  
slurries, thermal processing of cowpea 1993
- Cowpea flour**  
flow behaviour/gelatinization of cowpea flour 1994
- Cowpea starch**  
flow behaviour/gelatinization of cowpea starch 1994
- Crocodile**  
slaughter/dressing of crocodiles 2084
- Crystallization**  
parameters, on-line estimation of crystallization 2048
- Cucumber**  
acidification & microorganisms during fermentation of cucumber 2018
- Cucurbita**  
see Pumpkin
- Cumin**  
seed mycoflora of cumin 2153
- Dairy**  
gels, microstructure & compression strength of dairy 2064
- Dairy processing**  
biotechnology/separation technology application in dairy processing 2063
- Dairy product**  
bifidobacteria enumeration in dairy products 2068  
sensory quality optimisation RSM of dairy products 2070  
spoilage monitoring predictive microbiology of dairy products 2069
- Debitting**  
orange juices, processing & debittering of California Navel 2137
- Decay**  
grapefruit, thlabendazole/imazalil & decay of 2031
- Dehydration**  
prediction mathematical model for osmotic dehydration 1959
- Dietary fiber**  
components, legume processing effect on dietary fiber 1992
- Digestibility**  
weaning food, protein/starch digestibility of 2130
- Diphenylamine**  
apple, shelf-life extension diphenylamine in 2024
- Dried food**  
rice, spices/salt & storage stability of dehydrated 1984  
strain stress analysis of dehydrated foods 1961  
strawberries, processing/packaging/storage & quality of freeze-dried 2037
- Drumstick**  
thermal process for canned drumstick 2017
- Drying**  
egg powder, drying & properties of 2109  
figs, drying rates of solar dried 2030  
pumpkins, drying of 2020
- Egg**  
billed eggs, peeling properties of 2116  
niger cake-deoiled & production/quality of egg 2115  
pickled chicken eggs 2110  
quail eggs, storage stability of oil-based pickled 2114
- Egg albumen**  
antistherial effects of raw egg albumen 2111
- Egg powder**  
drying/packaging materials & properties of egg powder 2109  
lipid extraction & emulsifying properties/soluble protein fraction of egg yolk powder 2112
- Egg protein**  
egg yolk powder, lipid extraction & soluble protein fraction of 2112
- Egg yolk**  
phosvitin, emulsifying properties of egg yolk 2113
- Emulsifying capacity**  
corn germ protein flour, emulsifying capacity of 2082
- Emulsifying property**  
egg yolk phosvitin, emulsifying properties of 2113  
egg yolk powder, lipid extraction & emulsifying properties of 2112
- Emulsion**  
butter oil emulsions, *Aspergillus* niger lipase action on 2072  
Candelilla wax emulsion, preparation/stability of 1967  
corn germ protein flour, emulsion stability of 2082  
milk fat emulsions, bovine milk fat globule membrane & properties of 2067
- Energy**  
power reduction in condensing system 2045
- Enzyme**  
butter, fatty acid enzymatic analysis for 2071
- Equipment**  
evaporators, performance of falling film 2052  
sugar factory, entrainment prevention polybaffle entrainment catcher in 2044  
sugar industry, steam desuperheater exhaust in 2046
- Escherichia coli**  
finfish, *E. coli* on commercially processed Alaskan 2125
- Ester**  
sucrose carbonic acid ester, toxicological evaluation of 2157
- Esterification**  
Mucor miehei lipase, media & esterification catalysed by 1974
- Ethylene**  
tomatoes, ethylene & ripening of 2021
- Ethylmaleimide**  
lactoglobulin gels, ethylmaleimide & rheological properties of 1965
- Eugenol**  
antibacterial activity of eugenol 1976
- Evaporator**  
performance of falling film evaporators 2052
- Extraction**  
flavour, extraction supercritical CO<sub>2</sub> of 2152  
grape juice extraction technique standardisation 2032
- Extrudate**  
cassava extrudates, extrusion variables effect on 2013  
cassava flour extrudate, texture/microstructure of 2014
- Extrusion**  
cassava extrudates, extrusion variables effect on 2013  
soy flour, dry-extrusion expelling & production of medium fat 2006



- Faba bean**  
medamms, cooking temp./time & amino acid composition/antinutritional factors of cooked 1995
- Fat**  
rasogolla preparation, fat level effect on 2079  
soy flour, dry-extrusion expelling & production of medium fat 2006  
soy-wheat flour blend, storage stability of full-fat 2007
- Fats animal**  
beef/pork, fat & grinding time/pressure developed in 2102  
goat organ/muscle meat, fat in 2087
- Fats milk**  
emulsions, bovine milk fat globule membrane & properties of milk fat 2067  
modification of milk fat 2063
- Fats vegetable**  
triacylglycerol composition of vegetable fat 2141
- Fatty acid**  
butter, fatty acid flow injection/enzymatic analysis for 2071  
cheese flavours, fatty acids in aged Italian 2074  
fish, processing & fatty acid comp. in 2122  
potato, beef tallow-hydrogenated vegetable oil & volatile free fatty acids in French fried 2092  
safflower, fatty acid composition of 2002
- Fenvalerate**  
spectrophotometric detn. of fenvalerate 1969
- Fermentation**  
cucumber, acidification & microorganisms during fermentation of 2018
- Fermented food**  
meat spreads, pork mechanically deboned use in fermented 2107
- Fertilizer**  
sugarcane, fertilizers & yield/sucrose content of Bangladesh 2040
- Ficus carica**  
see Fig
- Fig**  
Ficus carica, chemical composition/drying rates of solar dried 2030
- Films**  
water vapour transmission rate of multi-layer flexible films 1958
- Finfish**  
Alaskan finfish, microorganisms on commercially processed 2125
- Fish**  
India, fish production prospects in 1956  
freshness detection, trimethylamine gas sensory for fish 2124  
processing & fatty acid comp. in fish 2122
- Fish oil**  
mayonnaise, quality of fish oil 2129
- Flavobacterium**  
finfish, Flavobacterium in commercially processed 2125
- Flavour**  
extraction supercritical CO<sub>2</sub> of flavour 2152  
meats, alkylphenols/thiophenol & flavour of red 2088  
wine/passion fruit juice, endo- $\beta$ -glucosidase & flavour of 2140
- Flavour compound**  
pineapple orange juice, lipid/composite film & volatile flavour compounds of 2136  
sorption LDPE film of flavour compounds 1968
- Flow**  
cowpea flour/starch, flow behaviour of 1994
- Flow-injection**  
butter, fatty acid flow injection analysis for 2071
- Fried food**  
potatoes, beef tallow-hydrogenated vegetable oil & volatile free fatty acids in French fried 2092
- Fries**  
quality assessment of industrial prefried French fries 2144
- Frozen food**  
pork patties, light/salt & oxidative stability of frozen 2106  
strawberries, processing/packaging/storage & quality of freeze-dried 2037
- Fruit**  
India, fruit production prospects in 1956
- Fruit juice**  
processing of fruit juices 2132
- Functional property**  
teary bean flour protein, functional properties of 1991
- Fungi**  
soy protein concentrate, fungal protease modification of 2008  
spices, seed mycoflora of 2153
- Gas chromatography**  
salmon, volatile compounds GC detn. in canned 2126
- Gel**  
beef protein fractions, gels heat-induced properties of 2101  
jam, fruit content & gel strength of strawberry/peach 2023  
microstructure & compression strength of dairy gels 2064  
surimi, water binding/ingredient dispersion pattern & gel texture of 2118
- Gelatinization**  
cowpea flour/starch, gelatinization of 1994
- Gellan gum**  
monovalent cations & isolation of gellan gum 1966
- Germination**  
soybean, storage & germination of 2004
- Globulin**  
Sesame indicum, chymotrypsin &  $\alpha$ -globulin of 2003  
teary bean globulin, functional properties of 1991
- Glucosidase**  
passion fruit juice/wine, endo- $\beta$ -glucosidase & flavour of 2140
- Glycol**  
lactoglobulin gels, propylene glycol & rheological properties of 1965
- Goat**  
meat, moisture/fat/cholesterol in goat organ/muscle 2087
- Gossypol**  
bovine serum albumin-gossypol interaction stoichiometric analysis 2086
- Grape**  
ripening, composition of kniffln trained grapes during 2033  
var., temperate condition & juice processing suitability of grape 2032
- Grape juice**  
grape var., temperate condition & juice processing suitability of 2032  
harvest date & quality of 'Perlette' grape juice 2135



- Grapefruit**  
thiabendazole/imazalil & chill  
injury/decay of grapefruit 2031
- Groundnut**  
harvest/maturity &  
properties/composition of  
groundnut 1998  
kernels, characterization of  
 $\gamma$ -irradiated stored  
groundnut 1997  
proteolytic enzymes & arachidins of  
groundnuts 1999
- Heat exchanger**  
particle residence time  
distribution & scraped surface  
heat exchanger 1962
- Heating**  
apple juice concentrate, heat  
treatment & physicochemical  
properties/microbial quality of  
2133  
inosine 5'-monophosphate, heating  
& sensory changes in 2155  
oysters, heating & quality of  
pasteurized 2119
- Homogenization**  
orange juice concentrate,  
homogenization & viscosity of  
2139
- Hydrogenation**  
soybean oil, hydrogenation of  
oxidized 2150
- Hydroperoxide**  
soybeans, hydroperoxide formation  
in stored 2005
- Imazalil**  
grapefruit, imazalil & chill  
injury/decay of 2031
- Industry**  
sugar factory, entrainment  
prevention polybaffle  
entrainment catcher in 2044  
sugar industry, steam  
desuperheater exhaust in 2046
- Inosine 5'-monophosphate**  
heating & sensory changes in  
inosine 5'-monophosphate 2155
- Iron**  
cereals, phytate & Fe  
availability in 1982
- Irradiated food**  
groundnut kernels,  
characterization of  
 $\gamma$ -irradiated stored 1997
- Irradiation**  
meat,  $\gamma$ -irradiation &  
colour/oxidative stability of  
nitrite-free cured 2090  
tomatoes,  $\gamma$ -irradiation &  
ripening of 2021
- Jam**  
pumpkin jam 2020  
strawberry jam, cooking &  
volatile compounds in 2038
- Kachhagolla**  
preparation of Kachhagolla 2078
- Kanaboko**  
axial compression properties of  
kanaboko 2128
- LDPE**  
flavour compounds, sorption LDPE  
film of 1968
- Lactobacillus casei**  
Cheddar cheese, L. casei &  
accelerated maturation of 2076
- Lactococcus lactis**  
citrate metabolism in L. lactis  
1973
- Lactoglobulin**  
gels, rheological properties of  
lactoglobulin 1965
- Lecithin**  
cheese, lecithin & yield/quality  
of soft 2073
- Legume**  
dietary fiber components, legume  
processing effect on 1992
- Lipase**  
Mucor miehei lipase & primary  
amines alkylation 1975  
Mucor miehei lipase, media &  
esterification catalysed by 1974  
butter oil emulsions, Aspergillus  
lipase action on 2072
- Lipid**  
avocado peels, ripening & lipid  
peroxidation products in 2028  
seal meat, lipid extraction from  
mechanically separated 2120
- Lipids**  
pineapple orange juice, lipid &  
volatile flavour components of  
2136
- Listeria**  
egg albumen, antilisterial  
effects of raw 2111
- Listeria monocytogenes**  
hydrostatic pressure & response  
of L. monocytogenes 1971
- Mackerel**  
processing & fatty acid comp. in  
mackerel 2122
- Malachite green**  
toxicity of malachite green 1981
- Mango**  
storage/ripening & quality  
changes in Tommy Atkins mangoes  
2034
- Marjoram**  
antioxidant activity of marjoram  
1977
- Maturation**  
Cheddar cheese, L. casei &  
accelerated maturation of 2076
- Maturity**  
groundnut, maturity &  
properties/composition of 1998
- Mayonnaise**  
fish oil mayonnaise, quality of  
2129
- Meat**  
alkylphenols/thiophenol & flavour  
of red meats 2088  
batters, air content detn. in  
meat 2091  
cholesterol oxidation products in  
raw meat 2108  
cured-meat pigment, synthesis of  
cooked 2089  
 $\gamma$ -irradiation &  
colour/oxidative stability of  
nitrite-free cured meat 2090  
India, meat production prospects  
in 1956  
Salmonella in meat 2117
- Mechanical property**  
groundnut, harvest date/maturity  
& mechanical properties of 1998  
soy protein gels, mechanical  
properties of heat-induced 2009
- Medammi**  
see Faba bean
- Menhaden oil**  
soybean oil/Menhaden oil blend,  
chemical/sensory properties of  
stored 2149
- Mercaptoethanol**  
lactoglobulin gels,  
mercaptoethanol & rheological  
properties of 1965
- Metabolism**  
Lactococcus lactis, citrate  
metabolism in 1973
- Metanil yellow**  
toxicity of metanil yellow 1981
- Methyl linoleate**  
 $a_w$  & autooxidation of stored  
methyl linoleate 1964  
oils, deuterium-labelled methyl  
linoleate from seed 2145
- Methylcellulose gum**  
cake, methylcellulose gum &  
structure/texture of  
microwave-baked 2062
- Microbial quality**  
apple juice concentrate,  
processing/heat treatment &  
microbial quality of 2133  
beef steaks, microbial quality of  
precooked restructured 2095



**Microbiology**  
dairy products, spoilage  
monitoring predictive  
microbiology of 2069

**Microorganism**  
cucumber, acidification &  
microorganisms during  
fermentation of 2018  
soybean, storage & microflora of  
2004

**Microscopy**  
carrot stick, scanning electron  
microscopy of 2012

**Microwave**  
canola seeds, microwave &  
myrosinase in 1996  
food, Na microwave digestion AAS  
analysis in 1970  
pepper, microwave & colour  
retention in 2154

**Milk**  
*Bacillus cereus* & quality of  
low-temp. stored milk 2065  
cow's milk, processing properties  
of crossbred 2066

**Milk product**  
Kachhagolla, preparation of 2078

**Milling**  
wheat flour, milling method &  
properties/breadmaking  
characteristics of 1987

**Moisture**  
goat organ/muscle meat, moisture  
in 2087

**Molasses**  
sugar colour & purity of molasses  
2054  
yeast distiller's & fermentation  
of molasses 2051

**Moraxella**  
finfish, *Moraxella* on  
commercially processed Alaskan  
2125

**Mucor miehei**  
lipase, media & esterification  
catalysed by *Mucor miehei* 1974  
lipase, primary amines alkylation  
& *Mucor miehei* 1975

**Muscle**  
goat muscle meat,  
moisture/fat/cholesterol in 2087

**Mustard oil**  
rice bran oil, rape-mustard oil  
detection in 2147

**Mutton**  
goat organ/muscle meat,  
moisture/fat/cholesterol in 2087

**Mycotoxin**  
rats, mycotoxins effect on  
pregnant albino 2158

**Myrosinase**  
canola seeds, microwave &  
myrosinase in 1996

**Nickel**  
vanaspatri, Ni content in 2142

**Nitrite**  
beef sausages, blood & nitrite  
antibotulinal efficacy in 2099  
meat,  $\gamma$ -irradiation &  
colour/oxidative stability of  
nitrite-free cured 2090

**Nitrogen**  
sugarcane, N<sub>2</sub> &  
growth/yield/quality of 2042

**Nutritional quality**  
faba bean medamms, cooking  
temp./time & antinutritional  
factors of cooked 1995

**Nutritional value**  
pumpkins, nutritional value of  
2020

**Oat**  
phytate & Fe availability in oat  
1982

**Oil**  
potatoes, beef  
tallow-hydrogenated vegetable  
oil & volatile free fatty acids  
in French fried 2092  
quality estimation of crude oil  
2143  
seed oil, deuterium-labelled  
methyl linoleate from 2145

**Okra**  
thermal process for canned okra  
2017

**Oncorhynchus keta**  
see Salmon

**Orange II**  
toxicity of orange II 1981

**Orange juice**  
Ca/processing & quality of orange  
juice 2138  
lipid/composite film & volatile  
flavour components of pineapple  
orange juice 2136  
processing & debittering of  
California Navel orange juices  
2137

**Orange juice concentrate**  
homogenization & viscosity of  
orange juice concentrate 2139

**Oregano**  
antioxidant activity of oregano  
1977

**Osmosis**  
dehydration, prediction  
mathematical model for osmotic  
1959

**Oxidation**

methyllinoleate,  $a_w$  &  
autooxidation of stored 1964  
soybean oil, hydrogenation of  
oxidized 2150

**Oyster**  
*Crassastrea virginica*, heating &  
quality of pasteurized 2119

**Packaging**  
Asparagus spears, packaging &  
quality changes in stored 2016  
strawberries, packaging & quality  
of freeze-dried 2037

**Packaging material**  
egg powder, drying & properties  
of 2109  
plum, wooden boxes & packaging of  
2036

**Paleness**  
veal, paleness of milk-fed  
refrigerated stored 2085

**Pan**  
automatic feed control in vacuum  
pans 2048

**Passion fruit juice**  
endo- $\beta$ -glucosidase & flavour  
of passion fruit juice 2140

**Peach**  
jam, fruit content & gel strength  
of peach 2023  
temp. & CO<sub>2</sub> injury/flesh  
softening in peaches 2035

**Pectin**  
citrus pectin, conformation of  
2029

**Peeling**  
boiled eggs, peeling properties  
of 2116

**Pepper**  
black pepper, seed mycoflora of  
2153  
microwave/blanching & colour  
retention in pepper 2154

**Peptides**  
2063

**Perilla oil**  
phospholipids antioxidant effect  
on perilla oil 1978

**Phenolic compounds**  
cheese, phenolic compounds &  
flavours in aged Italian 2074

**Phospholipid**  
perilla oil, phospholipid  
antioxidant effect on 1978

**Phosvitin**  
egg yolk phosvitin, emulsifying  
properties of 2113

**Physical property**  
beef steaks,  
tenderization/ageing/thawing &  
physical properties of 2093



- milk fat emulsions, bovine milk fat globule membrane & properties of 2067
- pork sausage, sodium lactate & physical properties of 2104
- Phytate**  
cereals, phytate & Fe availability in 1982
- Phytic acid**  
rapeseed protein, phytic acid & functional properties of detoxified 2000
- Pickle**  
chicken egg, pickled 2110  
pumpkin preserve 2020
- Pigment**  
meat pigment, synthesis of cooked cured 2089  
seal meat, pigment extraction from mechanically separated 2120
- Plum**  
wooden boxes & transportation of plum 2036
- Polyamine**  
apple, polyamine & softening inhibition in Golden Delicious/McIntosh 2026
- Polyphosphate**  
squid, polyphosphate & frozen storage of Indian 2121
- Pork**  
carnosine & rancidity inhibition of salted ground pork 2105  
meat spreads, pork mechanically deboned use in fermented 2107  
sausage, sodium lactate & shelf-life/properties of pork 2104  
temp./fat grinding time/pressure developed in pork 2102
- Pork patties**  
light/salt & oxidative stability of frozen pork 2106
- Potato**  
beef tallow-hydrogenated vegetable oil & volatile free fatty acids in French fried potatoes 2092  
thermal process for canned potatoes 2017
- Poultry**  
India, poultry production prospects in 1956
- Preserve**  
pumpkin preserve 2020
- Processing**  
apple juice concentrate, processing & physicochemical properties/microbial quality of 2133  
dietary fiber components, legume processing effect on 1992  
fish, processing & fatty acid comp. in 2122  
fruit juices, processing of 2132  
milk, processing properties of crossbred cow's 2066  
orange juices, processing & quality of 2138  
orange juices, processing & debittering of California Navel 2137  
strawberries, processing & quality of freeze-dried 2037
- Production**  
food production prospects in India 1956
- Protease**  
soy protein concentrate, protease modification of 2008
- Protein**  
weaning food, protein digestibility in malted 2130
- Proteins animal**  
beef protein fractions, gels heat-induced properties of 2101
- Proteins milk**  
emulsifying capacity/emulsion stability of milk proteins 2082  
milk protein hydrolysates 2063
- Proteolytic enzyme**  
groundnuts, proteolytic enzymes & arachin of 1999
- Pseudomonas**  
finfish, *Pseudomonas* on commercially processed Alaskan 2125
- Pulse**  
India, pulses production prospects in 1956
- Pumpkin**  
cucurbita, chemistry/technology of, review 2020
- Quality**  
Sorghum, protein quality of 1990  
cheese, lecithin & quality of soft 2073  
chum salmon, quality enhancement of 2127  
egg, niger cake-deoiled & quality of 2115  
fish oil mayonnaise, quality of 2129  
fries, quality assessment of industrial prefried French 2144  
grape juice, harvest date & quality of 'Perlette' 2135  
mangoes, storage/ripening & quality changes in Tommy Atkins 2034  
milk, *B. cereus* & quality of low-temp. stored 2065  
oil, quality estimation of crude 2143  
orange juices, Ca/processing & quality of 2138  
oysters, heating & quality of pasteurized 2119  
strawberries, processing/packaging/storage & quality of freeze-dried 2037  
sugarcane, N<sub>2</sub> & quality of 2042  
sugarcane, crop geometry & physiology/juice quality of 2043  
sugarcane, farm yard manure/castor cake/press mud cake & quality of 2041
- Radiation**  
oxic radiation damage control by caffeine, review 2131
- Rainbow trout**  
adenine nucleotide degradation in modified atm. chill-stored rainbow trout 2123
- Rancidity**  
pork, carnosine & rancidity inhibition of salted 2105
- Rapeseed oil**  
rice bran oil, rape-mustard oil detection in 2147
- Rapeseed protein**  
phytic acid & functional properties of detoxified rapeseed protein 2000
- Rasogolla**  
preparation, fat/malda level effect on rasogolla 2079
- Ready-to-serve beverage**  
sugarcane juice ready-to-serve 2049
- Response surface methodology**  
dairy products, sensory quality optimization RSM of 2070
- Rheological property**  
wheat flour, milling method & rheological properties of 1987
- Rhodamine B**  
toxicity of rhodamine 1981
- Rice**  
beef/wild rice mixtures, properties of ground 2098  
cement from rice husk-ash, review 1983  
dehydrated rice, spices/salt & storage stability of 1984  
stickiness, starch granule-associated protein on cooked rice 1985



**Rice bran oil**  
rape-mustard oil detection in  
rice bran oil 2147

**Rice protein**  
stickiness, starch  
granule-associated protein on  
cooked rice 1985

**Rice starch**  
stickiness, starch  
granule-associated protein on  
cooked rice 1985

**Ripening**  
avocado peels, ripening & lipid  
peroxidation products in 2028  
Cheddar cheese, spray drying for  
accelerated ripened 2077  
cheese ripening-accelerated 2063  
grapes during ripening,  
composition of knifflin trained  
2033  
mangoes, ripening & quality  
changes in Tommy Atkins 2034  
tomatoes,  
ethylene/ $\gamma$ -irradiation &  
ripening of 2021

**Roti**  
corn characteristics & roti  
quality 1988

**Rye**  
phytate & Fe availability in rye  
1982

**Saccharomyces cerevisiae**  
sugar beet pulp, Sacch.  
cerevisiae production from 2010

**Safflower**  
Carthamus tinctorius, fatty acid  
composition of 2002  
volatile components of safflower  
2001

**Salmon**  
Oncorhynchus keta, quality  
enhancement of chum 2127  
Oncorhynchus, volatile compounds  
GC detn. in canned 2126

**Salmonella**  
meat/seafoods, Salmonella in 2117

**Salt**  
pork patties, salt & oxidative  
stability of frozen 2106  
rice, salt & storage stability of  
dehydrated 1984 1984

**Sauce**  
pumpkin sauce 2020

**Sausage**  
beef sausage, blood & nitrite  
antibotulinal efficacy in 2099  
pork sausage, sodium lactate &  
shelf-life/properties of 2104

**Seafood**  
Salmonella in seafoods 2117

**Seal**  
Phoca groenlandica meat,  
lipid/pigment extraction from  
mechanically separated 2120

**Sensory property**  
beef steaks,  
tenderization/ageing/thawing &  
sensory properties of 2093  
Menhaden oil/soybean oil blend,  
chemical/sensory properties of  
stored 2149  
pork sausage, sodium lactate &  
sensory properties of 2104

**Sensory quality**  
dairy products, sensory quality  
optimization RSM of 2070

**Sesame**  
Sesamum indicum, chymotrypsin &  
 $\alpha$ -globulin of 2003

**Shelf-life**  
apple, shelf-life extension  
CaCl<sub>2</sub>/bavistin/  
diphenylamine in 2024  
pork sausage, sodium lactate &  
shelf-life of 2104

**Slaughtering**  
crocodiles, slaughtering of 2084

**Slurry**  
cowpea slurries, thermal  
processing of 1993

**Sodium**  
food, Na microwave digestion AAS  
analysis in 1970

**Sodium lactate**  
beef top rounds, sodium lactate &  
acceptability of cooked roast  
stored 2096  
pork sausage, sodium lactate &  
shelf-life/properties of 2104

**Solar drying**  
figs, drying rates of solar dried  
2030

**Sorghum protein**  
Sorghum bicolor, protein quality  
of 1990

**Soy flour**  
bread soy flour fortified 2006  
extrusion-dry & production of  
medium fat soy flour 2006  
storage stability of full-fat  
soy-wheat flour blend 2007

**Soy protein**  
gels, texture/mechanical  
properties of heat-induced soy  
protein 2009

**Soy protein concentrate**  
protease modification of soy  
protein concentrate 2008

**Soybean**  
Glycine max, storage &

germination/microflora of 2004  
hydroperoxide formation in stored  
soybeans 2005

**Soybean oil**  
hydrogenation of oxidized soybean  
oils 2150  
isotherms, evaluation of soybean  
oil 2148  
menhaden oil/soybean oil blend,  
chemical/sensory properties of  
stored 2149

**Spearmint**  
antioxidant activity of spearmint  
1977

**Spectrophotometry**  
fenvalerate, spectrophotometric  
detn. of 1969

**Spice**  
India, spices production  
prospects in 1956  
rice, spices & storage stability  
of dehydrated 1984  
seed mycoflora of spices 2153

**Spoilage**  
beef, bacteria/volatile compounds  
& spoilage of ground 2097  
dairy products, spoilage  
monitoring predictive  
microbiology of 2069

**Spray drying**  
Cheddar cheese, spray drying for  
accelerated ripened 2077

**Spread**  
meat spreads, pork mechanically  
deboned use in fermented 2107

**Squid**  
Loligo duvaucell, polyphosphate &  
frozen storage of Indian 2121

**Stability**  
corn germ protein flour, emulsion  
stability of 2082  
eggs, storage stability of  
oil-based pickled quail 2114  
meat,  $\gamma$ -irradiation &  
oxidative stability of  
nitrite-free cured 2090  
pork patties, light/salt &  
oxidative stability of frozen  
2106  
rice, spices/salt & storage  
stability of dehydrated 1984  
soy-wheat flour blend, storage  
stability of full-fat 2007  
sunflower oil, stability of  
refined stored 2151

**Starch**  
weaning food, starch  
digestibility in malted 2130

**Steel**  
stainless steel, corrosion



- prevention of 1963
- Storage**
- eggs, storage stability of oil-based pickled quail 2114
  - groundnut kernels, characterization of  $\gamma$ -irradiated stored 1997
  - methyllinoleate,  $a_w$  & autooxidation of stored 1964
  - pumpkin, storage of 2020
  - rice, spices/salt & storage stability of dehydrated 1984
  - soy-wheat flour blend, storage stability of full-fat 2007
  - soybean, storage & germination/microflora of 2004
  - soybeans, hydroperoxide formation in stored 2005
  - strawberries, storage & quality of freeze-dried 2037
  - sunflower oil, stability of refined stored 2151
- Storage controlled atmosphere**
- apples, storage controlled atm. & quality of Golden Delicious 2025
- Storage fish**
- white-fish/rainbow trout, adenine nucleotide degradation in modified atm. chill-stored 2123
- Storage frozen**
- squid, polyphosphate & frozen storage of Indian 2121
- Storage fruits**
- apple, respiration model for stored 2027
  - mangoes, storage & quality changes in Tommy Atkins 2034
- Storage modified atmosphere**
- apples, storage modified atm. & quality of Golden Delicious 2025
- Storage vegetables**
- Asparagus spears, packaging & quality changes in stored 2016
  - Brassica, chlorophyll loss during storage of 2011
  - turnips, chemical composition of low-temp. stored 2015
- Strawberry**
- jam, cooking & volatile compounds in strawberry 2033
  - jam, fruit content & gel strength of strawberry 2023
  - processing/packaging/storage & quality of freeze-dried strawberries 2037
- Sugar**
- Bacillus subtilis* spores, caramelized sugar & germination response of 2059
  - factory, entrainment prevention polybaffle entrainment catcher in sugar 2044
  - factory, loss in sugar 2058
  - industry, condensate water treatment in sugar 2047
  - industry, steam & superheater exhaust in sugar 2046
  - molasses, sugarcane & purity of 2054
  - recovery of sugar 2057
- Sugar beet**
- pulp, *Sacch. cerevisiae* production from sugar beet 2010
- Sugarcane**
- crop geometry & physiology/juice quality of sugarcane 2043
  - farmyard manure/castor cake/press mud cake & yield/quality of sugarcane 2041
  - fertilizers & yield/sucrose content of Bangladesh sugarcane 2040
  - $N_2$  & growth/yield/quality of sugarcane 2042
- Sugarcane juice**
- crop geometry & juice quality of sugarcane 2043
  - microprocessor based control system for mixed juice flow 2050
  - ready-to-serve sugarcane juice 2049
  - sulphitation automatic of sugarcane juice 2056
  - sulphitation plants & clarification of sugarcane juice 2053
  - sulphiter & pH of sugarcane juice 2055
- Sulphitation**
- sugarcane juice, sulphitation automatic of 2056
  - sugarcane juice, sulphitation plants & clarification of 2053
- Sulphiter**
- sugarcane juice, sulphiter & pH of 2055
- Sunflower oil**
- stability of refined stored sunflower oil 2151
- Surimi**
- water binding/ingredient dispersion pattern & gel texture of surimi 2118
- Taste**
- corn germ protein flour products, phenolic acid & taste of 1989
- Technology**
- pumpkin, technology of, review 2020
- Temperature**
- kinetic parameter evaluation with linearly increasing temp. profile 2156
- Tenderization**
- beef steaks, tenderization & properties of 2093
- Tenderness**
- beef tenderness, prediction of 2094
- Tepary bean flour**
- protein, functional properties of tepary bean flour 1991
- Texture**
- cake, methylcellulose gum & structure/texture of microwave baked 2062
  - cassava flour extrudate, texture of 2014
  - soy protein gels, texture of heat-induced 2009
  - surlimi, water binding/ingredient dispersion pattern & gel texture of 2118
- Thawing**
- beef steaks, thawing & properties of 2093
- Thermal process**
- conduction-heated foods, thermal process lethality variability in 1960
  - cowpea slurries, thermal processing of 1993
  - vegetables, thermal process for canned 2017
- Thermal property**
- whey protein concentrates, thermal properties & solubility of 2081
- Thiabendazole**
- grapefruit, thiabendazole/imazalil & chill injury/decay of 2031
- Thiophenol**
- meats, thiophenol & flavour of red 2088
- Thyme**
- antioxidant activity of thyme 1977
- Tin**
- sunflower oil, stability of refined stored 2151
- Tomato**
- chilling & surface colour changes in tomatoes 2022
  - ethylene/ $\gamma$ -irradiation & ripening of tomatoes 2021
- Toxicity**
- colours, toxicological evaluation of non-permitted food 1981
  - sucrose carbonic acid ester,



toxicological evaluation of 2157

**Transportation**

plum, wooden boxes &  
transportation of 2036

**Triacylglycerol**

vegetable fat, triacylglycerol  
composition of 2141

**Trimethylamine**

fish freshness detection,  
trimethylamine gas sensory for  
2124

**Turnip**

chemical composition of low-temp.  
stored turnips 2015

**Vanaspati**

NI content in vanaspati 2142

**Veal**

paleness of milk-fed refrigerated  
stored veal 2085

**Vegetable**

amaranth, protein fractions of  
grain 2019

India, vegetable production  
prospects in 1956

**Vibrio parahaemolyticus**

hydrostatic pressure & response  
of parahaemolyticus 1971

**Viscosity**

cottonseed oil, viscosity of 2146

orange juice concentrate,  
homogenization & viscosity of  
2139

**Volatile compound**

beef, volatile compounds &  
spoilage of ground 2097

safflower, volatile components of  
2001

salmon, volatile GC detn. in  
canned 2126

**Volatile compounds**

strawberry jam, cooking &  
volatile compounds in 2038

**Waste water**

agro-based industries effluents,  
useful materials from 1957

condensate water treatment in  
sugar industry 2047

**Water**

surimi, water binding & gel  
texture of 2118

**Weaning food**

protein/starch digestibility in  
malted weaning food 2130

**Wheat**

physico-chemical characteristics  
of Indian wheat 1986

phytate & Fe availability in  
wheat 1982

**Wheat flour**

malda level effect on rasogolla  
preparation 2079

milling method &

properties/breadmaking

characteristics of wheat flour  
1987

storage stability of full-fat

soy-wheat flour blend 2007

**Whey**

acid whey, antioxidant from 2080

**Whey protein concentrate**

thermal properties & solubility  
of whey protein concentrates 208

**White fish**

adenine nucleotide degradation in  
modified atm. chill stored white  
fish 2123

**Wine**

endo- $\beta$ -glucosidase & flavour  
of wine 2140

**Yam**

elephant yam, thermal process for  
canned 2017

**Yeast**

yeast distiller's & fermentation  
of molasses 2051



**NATIONAL INFORMATION CENTRE FOR FOOD SCIENCE AND TECHNOLOGY  
CFTRI, MYSORE - 570 013**

*Also subscribe to our other periodicals*

**1. FOOD DIGEST (Quarterly)**

This is oriented towards the information needs of food industries, trade and marketing personnel, entrepreneurs, decision makers and individuals engaged in food field.

**Annl Subn: Indian Rs. 150/- Foreign \$. 65/-**

**2. FOOD PATENTS (Quarterly)**

Gives world patent information on Food Science and Technology taken from national and international sources. Abstracts are provided for food patents registered in India.

**Annl Subn: Indian Rs. 100/- Foreign \$. 50/-**

*For details, please write to:*

*The Area Co-ordinator*

*FOSTIS, CFTRI*

*Mysore - 570 013, India*

**COMPUTERISED DATABASE SEARCH OF WORLD FOOD LITERATURE**

The National Information Centre for Food Science and Technology (NICFOS) at CFTRI, Mysore, has developed facilities for computerised "Database Search" of the world literature in Food Science and Technology and related disciplines. Retrospective search-service facility extending upto about 20 years is available on nominal payment basis. Titles alone or with abstracts can be requisitioned as per the needs of the intending users.

*For details, please write to:*

*The Area Co-ordinator*

*FOSTIS, CFTRI*

*Mysore - 570 013, India*



Regd. No. 36729/84